

DANGEROUS POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
CAN KILL IF SWALLOWED
DO NOT PUT IN DRINK BOTTLES
KEEP LOCKED UP**

EzyCrop Paraquat 250 SL Herbicide



**RLP
Approved**



**ACTIVE CONSTITUENT: 250 g/L PARAQUAT
present as PARAQUAT DICHLORIDE**

GROUP	L	HERBICIDE
--------------	----------	------------------

**EzyCrop Paraquat 250 SL Herbicide is for the control of a
wide range of Grasses and Broadleaf Weeds as per
Directions for Use**

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

**CONTENTS: 20 LITRES
(5 L, 110L, 200L)**

EzyCrop Pty Ltd
2/22 Horne Street
Elsternwick
VIC 3185
Australia

Tel: 03 9505 0044
ACN: 156 476 827

DO NOT USE THIS PRODUCT IN THE HOME GARDEN.**STORAGE AND DISPOSAL****(5, 20 and 200L only)**

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For Refillable Containers (110L)

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Very dangerous, particularly the concentrate. Product is poisonous if swallowed. Will irritate the nose, throat and skin. Attacks the eyes, protect the eyes while using. Avoid contact with the eyes, skin and clothing. When opening the container and preparing product for use wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. **DO NOT** inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

SPRAY APPLICATION

Do not work in spray mist.

Do not continue to use if skin irritation or nose-bleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.

When there is a risk of exposure to spray mist wear waterproof footwear and waterproof protective clothing, impervious gauntlet length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended but in any event use a respirator that complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.

Avoid contacting vegetation wet with spray but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

FIRST AID

If poisoning occurs get to a doctor or hospital quickly. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet that can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of the EzyCrop Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. EzyCrop Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. EzyCrop Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

UN No. 3016	BIPYRIDILIUM PESTICIDE LIQUID, TOXIC, (contains paraquat)
In a Transport Emergency dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA-WIDE
PG III	HAZCHEM 2 X

TOXIC 6

APVMA Approval No: 67977/59893

Batch No.
DOM

DANGEROUS POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
CAN KILL IF SWALLOWED
DO NOT PUT IN DRINK BOTTLES
KEEP LOCKED UP**

EzyCrop Paraquat 250 SL Herbicide

**ACTIVE CONSTITUENT: 250 g/L PARAQUAT
present as PARAQUAT DICHLORIDE**

GROUP	L	HERBICIDE
--------------	----------	------------------

**EzyCrop Paraquat 250 SL Herbicide is for the control of a
wide range of Grasses and Broadleaf Weeds as per
Directions for Use**

IMPORTANT: READ THIS LEAFLET BEFORE USE

APVMA Approval No: 67977/59893

EzyCrop Pty Ltd
2/22 Home Street
Elsternwick
VIC 3185
Australia

Tel: 03 9505 0044
ACN: 156 476 827

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT add wetter unless spraying at high volume. Where EzyCrop Paraquat 250 SL is mixed with water at less than 400mL/100L of water, add 100mL Wetter 1000 per 100L of spray.

DO NOT spray plants which are waterlogged, under stress of any kind or covered with soil or dust.

DO NOT spray plants covered with heavy dew, but rain following spraying will not affect results.

DO NOT sow or cultivate for 1 hour after spraying but operations should commence within 7 days.

For ground application only – do not use through aircraft, misting machines or hand-held ultra low volume controlled droplet applications (CDA units).

Crop Use or Situation	Weeds Controlled	State	Rate/ha	Critical Comments
Aid to Cultivation to minimise cultivation and prepare a clean bed for sowing.	Annual grass and broadleaf weed control.	Qld, NSW, Vic, SA, Tas, NT, ACT only	1.2 to 1.6L	Where cultivation follows spraying, it may commence one hour after spraying but should be completed within 7 days. Where heavy weed growth is present at spraying a better seedbed will result if cultivation is delayed 3-5 days.
	Early autumn sowing.		1.6 to 2.4L	Use the higher rates for dense, more mature weed stands. Wild oats must have at least two leaves. Where "Reglone" is used the lower EzyCrop Paraquat 250 SL rate should be sufficient to control dense, mature weeds.
	Winter, spring and early summer sowing.	Qld, Vic, SA, Tas, NT only	600 to 800 mL	Pasture: Remains of old pasture should be reduced by continuous heavy grazing. Remove stock 3-5 days before spraying to allow to freshen up.
	Wild oats at 2-5 leaf stage in autumn / winter.		600 mL	
Rice	Annual grass and broadleaf weed control.	Qld, NSW, NT only	1.6L	Pre-sowing.
			800 mL	Post-sowing, pre-crop emergence.
Wild Oat control in Spring Fallows	Wild oats at 2 to 5 leaf stage.	Qld, NSW, NT, ACT only	1.2 to 2 L	Use higher rate for summer growth. Avoid spraying under hot, dry conditions. Best results will be obtained when spraying is carried out in the late evening.
Kikuyu / Paspalum Pasture	To suppress growth to oversow winter seed.	Qld, NSW, ACT only	1.6 or 2.4 L	Use the high rate for February spraying and the low rate in March.
Selective Weed Control Autumn / early Winter annual and perennial clover	Annual grass and some broadleaf weed control except Paterson's Curse, Sorrel, Dock, Shepherd's Purse and some thistles.	All States	600mL to 1.2L	Use the higher rate for dense weed stands
	Control of some broadleaf weeds, including; Patterson's Curse, Sorrel, Dock, Shepherd's Purse and some thistles will not be achieved. Alternative methods such as the spray-graze technique with 2,4-D or MCPA should be considered.		1.2 to 1.6L	
Late winter/early spring - Annual and perennial clovers - Cocksfoot - Perennial ryegrass - Phalaris - Demeter fescue only		Qld, NSW, Vic, SA, Tas, NT, ACT only	1.6 to 2.4L	Use the higher rate in winter/early spring when barley grass is present. All applications: Graze pastures continuously after the seasonal break to a height of 2-4 cm. Remove stock 2-3 days before spraying to allow weeds to freshen up. Do not apply until clover has reached the 6 leaf stage. Do not spray clovers, which are affected by insect attack, disease or moisture stress. Do not use on clover pastures growing in water repellent sands or other situations subject to moisture stress at or immediately following treatment. Poor recovery of the clover will result. Use the lower rate for Cocksfoot and Perennial Ryegrass and the higher rate for

Crop Use or Situation	Weeds Controlled	State	Rate/ha	Critical Comments
				Phalaris and Demeter Fescue. The perennial grasses must be at least 12 months old at spraying. DO NOT APPLY TO MEDICS
	Yorkshire Fog Grass		1.2 L	Apply in early spring to reduce Yorkshire Fog Grass component and increase the clover and desirable grass component. Mixed pastures will be scorched initially but should show good recovery and beneficial changes in composition following spring rainfall and growth. In lower rainfall areas application in mid to late winter may be almost as effective but allow better pasture recovery. If pasture has been grazed allow for sufficient time for pasture and fog grass recovery before spraying. Apply in spray volumes of 100 to 250 L/ha, the latter for dense or tall ungrazed pastures. Add Wetter 1000 at 120ml per 100L.
Lucerne Autumn/ early winter	Annual grass and some broadleaf weeds.	Qld, Vic, SA, WA, Tas, NT only	1.2 to 1.6L	Use the higher rates for dense weed stands. Do not spray Lucerne stands under 12 months old. If mintweed is present use atrazine (900g/kg) at 600g/ha.
		NSW only	1.2L	WARNING - In certain areas, an uncommon species of barley grass (<i>H. glaucum</i> - common species of barley grass is <i>H. leporinum</i>) resistant to paraquat based products has become established. It may regrow after an initial scorch by EzyCrop Paraquat 250 SL. Where this problem is suspected use fluazifop-p-butyl for grass weed control. If EzyCrop Paraquat 250 SL has been applied use fluazifop-p-butyl at 1L/ha after regrowth but before heading.
Late winter/early Spring		Qld, Vic, SA, WA, Tas, NT only	1.6 to 2.4L	
		NSW, ACT only	1.2L	
Perennial Grass Seed Crops Cocksfoot, Perennial ryegrass, Phalaris and Demeter Fescue only	Annual grass and some broadleaf weeds.	All States	600mL to 1.2L	Use the low rate for Cocksfoot and perennial ryegrass and the higher rate for Phalaris and Demeter Fescue. Spray about 4 weeks after a full weed germination following the autumn break. The perennial grasses must be at least 12 months old at spraying.
Spray topping to reduce seed set Chickpeas Faba beans Field peas Lentils Lupins Vetch	Annual ryegrass	All States	400mL or 800mL	As an aid in managing annual ryegrass resistance. For use on escapes from a previous herbicide application in the current crop. Spray the crop when the ryegrass is at the optimum stage, that is when the last ryegrass seed heads at the bottom of the plant have emerged and the majority are at or just past flowering (with anthers present or glumes open) but before haying-off is evident - usually October to November. Use of the higher rate in these crops is usually more reliable and gives a greater reduction in seed set. Reduction in crop yield may occur especially if the crop is less advanced relative to the ryegrass - that is if crops have a majority of green immature pods. The higher rate may also increase any yield reduction.

Crop Use or Situation	Weeds Controlled	State	Rate/ha	Critical Comments
				In practice crop losses in excess of 25% may occur. Apply by ground boom only in 50-100L/ha. Spray with a calibrated boom spray raised to give double overlap at the level of the ryegrass seed heads. Pressures of 250 -350 kPa and use of 110015 or 02 nozzles or equivalent will aid coverage.
Spray topping to reduce seed set Pastures	Grasses generally (particularly annual ryegrass).	All States	400mL	Heavily graze paddocks during spring flush to encourage even head development. Remove stock 2-3 weeks before the anticipated maturity date of the target species. However, if this is not feasible through lack of stock it is preferable to allow the pasture to mature ungrazed. Delay spraying until the last seed-heads at the bottom of the plant have emerged and initial signs of haying-off appear. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.
	Barley grass			Manage paddocks as above. Spray after head emergence but when all seed heads are green and there is no sign of haying-off. Inspect paddocks before returning stock. Provided spraying was carried out before hardening seeds are present harrow to knock seed from the heads. Do not introduce lambs into paddock until safe from risk of seed injury. If seasonal conditions favour regeneration, stock should be returned to selectively graze new shoots. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.
	Saffron thistle	NSW, SA, ACT only		Spray after the plant begins to run to head until flowering.
Prevention of annual ryegrass toxicity	Spray top - graze to destroy seed heads.	WA only	400mL	Grazing management as for spray topping above. Remove stock 3-4 weeks before anticipated maturity date. Spray must be applied within 10 days after emergence of the first ryegrass seed heads. To ensure adequate control of toxin development, <u>heavy continuous grazing is essential from 1 day after spraying</u> until the pasture has completely hayed off. The required stocking rate will vary but must be sufficient to keep all regrowth after spraying completely eaten off to prevent further growth producing new seed heads, which could become toxic.
Hay Freezing	Maximum retention of protein in standing dry feed.	All States	800mL	Graze paddocks as for spray topping above. Remove 3-4 weeks before the anticipated maturity date. Apply prior to commencement of haying-off regardless of the grass species involved. Spray with a calibrated boom spray raised to give double overlap at the level of the seed heads.
General Weed control	Annual weed control	Qld, NSW, NT	1.6 to 3.2 L sprayed ha.	Spray as necessary for control of annual weeds. Avoid contacting crop foliage.

Crop Use or Situation	Weeds Controlled	State	Rate/ha	Critical Comments
Bananas		only	160 to 320 ml per 100L Misters 8mL/L	Apply soon after emergence and before weeds reach 15cm in height. Use spraying pressure less than 240kPa. Avoid chemical contact with roots and peepers near pseudo stem. Repeat as required.
Hops	Annual grasses	Vic, Tas, only	* 1.2 to 1.6L plus 1.1 kg/ha simazine (900g/kg) and/or 750mL to 1.4L/ha diquat 200g/L	Apply as a directed inter-row spray prior to crop emergence from winter dormancy, using a minimum of 250 L/ha spray volume to ensure good and even coverage of weeds.
Orchards and Vineyards	Annual weed control	Qld, Vic, SA, WA, Tas, NT only	* + 1.6 to 3.2 L per sprayed ha (a) (b) see below	Spray as necessary for control of annual weeds. Avoid contacting crop foliage. EzyCrop Paraquat 250 SL will not harm trees or vines with mature brown bark if this alone is sprayed. Use the higher rate for dense weed growth.
		NSW only	* + 1.7L per sprayed ha (a) (b) see below	If fat hen <i>Chenopodium album</i> or <i>Portulaca</i> spp. are present and EzyCrop Paraquat 250 SL rate is less than the ratio 800mL/100L add 120mL 1000g/L non-ionic wetter per 100L of spray mix.
Peanuts Post-emergence (in-crop)	<i>Datura</i> spp. (2-4 leaf)	Qld, NT only	400mL	Spray peanuts up to 7-8 leaf stage but before majority of plants flowering. Foliage will be scorched following application but plants recover rapidly. Apply in 200-250 L/ha for thorough coverage of weed foliage. A dense canopy of weeds may reduce weed control due to shielding. Add 60mL 1000g/L non-ionic wetter per 100L of spray mix.
	Annual ground cherry (2-3 leaf)		600mL	
	Apple-of-Peru (2-4 leaf)			
	Milkweed (2-3 leaf)			
	Stagger weed (2-3 leaf)		800mL	
	Blue heliotrope (2-3 leaf)			
Potatoes	General weed control (in-crop)	All States	* 1.2 to 1.6L	Spray at early crop emergence (no later than 25% emergence of potato shoots). Use the higher rate for dense weed growth.
	Pre-harvest weed control		* 2.8L	Spray about one week before digging and after tops have died down.
Row Crops, Vegetables and Market Gardens	Pre-planting and pre-crop emergence	All States	* + 1.2 to 1.6L or 200mL/100L	To control weeds in seedbeds. Treat no less than three days before sowing or before crop emergence. Use the lower rate for early autumn applications.
	Post-emergence inter-row weed control		* + 1.2 to 1.6L or 200mL/ 100L	Apply after crop seedlings have emerged or when transplanted crops are established. Direct the spray so that it does not touch the crop. Use shielded nozzles.
	Seedling weeds			Seedling weeds - use the lower rate for early

Crop Use or Situation	Weeds Controlled	State	Rate/ha	Critical Comments
				autumn applications.
	Older weeds		* 2.4L or 400mL/100L	More mature stages of weed growth.
Sugar Cane (Plant and ratoon)	Grass and some broadleaf weeds	Qld, NSW, NT only	+ 1.2 to 1.6L per sprayed ha	<p>Apply as a broadcast spray over-the-top of plant cane up to the 3-4 leaf stage. Cane foliage will be scorched but new leaves will appear in 7-10 days.</p> <p>Between the 3 to 4 leaf stage and the formation of the true stem use a directed, interspace spray with droppers and/or shields or leaf deflectors to avoid excessive spray drift onto cane foliage while spraying up to the cane bases. Use coarse nozzles such as flood jets (reflex nozzles) and pressure of 100-200 kPa. After the formation of the true stem, which is resistant to EzyCrop Paraquat 250 SL, droppers can be raised to overlap the spray pattern to give weed control in the stool. Use the higher rate for dense more mature weeds.</p> <p>EzyCrop Paraquat 250 SL can be mixed with EzyCrop Atrazine 900WG to give residual weed control when used as a blanket or directed spray (refer to atrazine label for specific rates).</p>
Non-Agricultural situations, around sheds, roadways, paths	Annual weed control	All States	* + 1.6 to 4 L/ha or 200mL/100L	<p>Spray to thoroughly wet weed growth. EzyCrop Paraquat 250 SL can be combined with soil residual herbicides EzyCrop Simazine 900WG or EzyCrop Atrazine 900WG to give rapid knockdown and prolonged weed control. Use the higher rate for dense weed growth.</p>
	Columbus grass	NSW only	+ Spot spraying 160mL/ 100L plus 1L flupropanate (745g/L) Boomspray 2.3 to 4.5L/ha plus 12 to 22L flupropanate (745g/L)	
Firebreaks	Knock down weed growth to eliminate fire hazard or assist firebreak burn	All States	1.6 to 4L	<p>Apply mid-winter to early summer.</p> <p>Use the higher rate for dense weed growth. After desiccation is complete the sprayed area may be burnt (normally 7–10 days after spraying).</p> <p>EzyCrop Paraquat 250 SL can be combined with soil residual herbicides EzyCrop Atrazine 900WG or EzyCrop Simazine 900WG to give rapid knockdown and prolonged weed control.</p>

* Capeweed or Erodium spp. present: Add diquat (200g/L) at 750mL to 1.5L/ha (125mL to 250 mL/100L for high volume spraying). Use higher rate for plants more than 10 cm diameter.

+ If EzyCrop Paraquat 250 SL Herbicide rate is less than the ratio 400mL/100L add 60mL 1000g/L non-ionic wetter per 100L of spray mix.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

WITHHOLDING PERIOD

DO NOT GRAZE OR CUT SPRAYED VEGETATION FOR STOCK FOOD FOR AT LEAST 1 DAY, OR GRAZE HORSES FOR 7 DAYS AFTER APPLICATION.

REMOVE STOCK FROM TREATED AREAS 3 DAYS BEFORE SLAUGHTER.

CHICKPEAS, FABA BEANS, FIELD PEAS, LENTILS, LUPINS – DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

PLEASE NOTE EXTRA WETTER REQUIREMENTS FOR HIGH VOLUME SPRAYING.

GENERAL INSTRUCTIONS**RESISTANT WEEDS WARNING**

GROUP	L	HERBICIDE
-------	---	-----------

EzyCrop Paraquat 250 SL Herbicide is a member of the bipyridyl group of herbicides. EzyCrop Paraquat 250 SL has the inhibitor of photosynthesis at Photosystem I mode of action. For weed resistance management EzyCrop Paraquat 250 SL is a Group L herbicide.

Some naturally occurring weed biotypes resistant to EzyCrop Paraquat 250 SL and other Group L herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by EzyCrop Paraquat 250 SL or other Group L herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, EzyCrop Pty Ltd accepts no liability for any losses that may result from the failure of EzyCrop Paraquat 250 SL to control resistant weeds.

This product kills annual grasses and most annual broadleaf weeds (excluding capeweed) in specified situations and should not be used for any other purpose. Quickly kills green plant tissue on contact. Is immediately inactivated in the soil or heavy dew. The principle of selective weed control with this product is that annual weeds are killed but perennial plants and clovers recover after an initial scorch. The control of annual weeds by spraying with this product will allow the desirable perennial species to thicken up at the expense of the weeds. Moisture and fertility should not be limiting at spraying and the proportion of desirable species must be great enough for them to fill in the areas previously occupied by weeds. Long-term weed control can be obtained following the quick knockdown given by this product if it is combined with soil residual chemicals.

- 1 Do not use hand-held ultra low volume controlled droplet applicators (CDA units), boomless jets or misting machines (except in banana plantations).
- 2 **Mixing**
Add the required quantity of product to water in the spray tank and agitate to give even mixing. Agitate again if left standing.
- 3 **Wetting agent**
This product contains a wetting agent and additional wetter is not required unless high volume spraying results in excessive dilution of wetter content. This will occur when product rates fall below 400mL per 100L of spray. Under such circumstances wetter should be added at the rate of 100mL of 1000g/L non-ionic wetter per 100L of spray mix.
Where Fat Hen or Portulaca are present in orchard or vineyard situations, extra wetter should be used when this product ratio is less than 800mL per 100L. Add wetter at double the above recommendations. Do not use alkaline or anionic wetting agents.
- 4 **Clean water**
Mix this product with **clean water** only. Water should be clean and free from clay, silt and algae. Providing it meets this requirement, saline water, water collected from roofs, bore water, dam water and water from creeks may be used.
- 5 **Application**
 - (i) **Cereals and Broadacre Spraying**
Use only through a properly calibrated boom spray that should be fitted with flat fan jets and adjusted to a height to give at least double overlap of the spray at the top of the weeds being sprayed. Spraying pressures should be in the range of 200-300kPa. Speed of travel should be

in the range of 6 – 15km/hr. It is essential that a good marking system be used. If a disc marker is used, it must be mounted so as to turn the soil back on to the area sprayed. It is essential to obtain good leaf coverage with the spray and volumes of dilute spray must be adjusted according to density of weed growth. 100L/ha may be used for seedlings or well grazed weeds up to 2 cm high. For plant height 2-5cm use 150L/ha and up to 6-10cm use 200L/ha. Spray volumes may be as low as 50L/ha (30L/ha in WA) for weed growth below 5cm high, or for spray topping and hay freezing. Equipment must be appropriate to this volume, properly calibrated and fitted with spraying tips designed to give droplets in the 200-250 µm Volume Median Diameter range.

(ii) **High Volume Application**

Higher volumes will generally be required to give good coverage of weed growth in situations other than those specified under cereals and other broadacre crops.

- (iii) Wash spray equipment with clean water immediately after use. This product is highly corrosive to metals, particularly galvanised iron and aluminium and should not be left for long periods in tanks or equipment made of these materials.

6 Compatibility

This product combines satisfactorily with the soil active herbicides EzyCrop Atrazine 900WG and EzyCrop Simazine 900WG where prolonged weed control is required as well as a quick knockdown. This product is compatible with diquat, dicamba, dicamba + MCPA, MCPA Amine (no more than 1L per 800mL EzyCrop Paraquat 250 SL Herbicide), chlorsulfuron, tri-allate and trifluralin.

7 Spraying conditions

Avoid spraying plants under stress from waterlogging, frost, drought etc. or covered with dust and soil. Results will be better if application is made in dull weather or at the end of the day. Light rain following spraying will not affect results. Avoid drift into neighbouring crops.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto susceptible plants/crops, cropping lands or pastures. This formulation should not be applied on or near water that is used for irrigation purposes.

PROTECTION OF LIVESTOCK

Domestic pets and poultry – keep away from treated areas.

This formulation should not be applied on or near water, which is used for livestock watering.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container. This formulation should not be applied on or near water, which is used for human consumption, livestock watering or irrigation purposes, or water used for commercial or recreational fishing.

STORAGE AND DISPOSAL

(5, 20 & 200L ONLY)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers.

Triple or (preferably) pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For Refillable Containers (110L)

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

GENERAL

Very dangerous, particularly the concentrate. Product is poisonous if swallowed. Will irritate the nose, throat and skin. Attacks the eyes, protect the eyes while using. Avoid contact with the eyes, skin and clothing. When opening the container and preparing product for use wear elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If clothing becomes contaminated with product remove clothing immediately. If product in eyes, wash it out immediately with water. Avoid contact with spray mist. **DO NOT** inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

SPRAY APPLICATION

- Do not work in spray mist.
- Do not continue to use if skin irritation or nose bleed occurs. This may be caused by exposure to spray mist as the result of incorrect use of equipment or adverse climatic conditions. Stop and review handling and spraying techniques before further spraying. If symptoms persist seek medical advice.
- Where there is a risk of exposure to spray mist, wear waterproof footwear and waterproof protective clothing, impervious gauntlet length gloves (rubber or PVC), goggles and a face mask and respirator covering nose and mouth and capable of filtering spray droplets. A high efficiency type particulate respirator is recommended but in any event use a respirator that complies with the requirements of AS1716 (Standards Association of Australia). Further advice on safety equipment should be obtained from a safety equipment manufacturer.
- Avoid contacting vegetation wet with spray but if necessary to do so wear waterproof footwear and waterproof protective clothing and gloves.

FIRST AID

If poisoning occurs get to a doctor or hospital quickly. Phone Australia 13 11 26. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet that can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of the EzyCrop Pty Ltd. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. EzyCrop Pty Ltd accepts no responsibility for any consequences resulting from the use of this product. EzyCrop Pty Ltd will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

In a Transport Emergency Dial 000 Police or Fire Brigade

UN No. 3016	BIPYRIDILIUM PESTICIDE LIQUID, TOXIC, (contains paraquat)
In a Transport Emergency dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA-WIDE
PG III	HAZCHEM 2 X

<p style="text-align: center;">TOXIC 6</p>
