

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

EzyCrop Propyzamide 500 SC Herbicide



**ACTIVE CONSTITUENT: 500 g/L PROPYZAMIDE
SOLVENT: 42 g/L ETHYLENE GLYCOL**

GROUP	D	HERBICIDE
--------------	----------	------------------

**For selective control of certain grasses and broad leaf
weeds in Lettuce, Sports Turf, Home Lawns and Legumes
Seed Crops and Pastures, as per the Direction for Use**

**Contents: 5L
(10L, 20L)**

EzyCrop Pty Ltd
2/22 Horne Street
Elsternwick
VIC 3185
Australia

Tel: 03 9505 0044
ACN: 156 476 827

DIRECTIONS FOR USE

Situation	Weeds Controlled	State	Rate L/ha	Critical Comments	
Lettuce Direct sown	Banyard Grass, Blackberry Nightshade, Chickweed,	All States	4.5 in 330 – 550L	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.	
Transplanted	Portulaca, Winter Grass, Ryegrass, Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles			Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4 cm of soil should be applied within 24 hours.	
		Vic only	2.3 in 300 – 550L	Apply on light sandy soils, low in organic matter.	
Turf & Lawns	Winter grass			Rate Bowling Green 37 x 37m	
Common Couch		All States	1.2 (12mL /100m ²)	170mL	Apply during the autumn and Winter, if established turf is infested with seedling winter grass. Ensure a uniform coverage. Irrigate immediately after application with 12-25 mm irrigation or rain and again 24 hours later. Repeat if necessary on Qld. Blue Couch after 4 weeks. DO NOT re-seed for 60 days after treatment. Use ONLY on established turf. DO NOT use on Bluegrass, ryegrass, fescue or bent grass lawns.
Queensland Blue couch		Qld only	0.6 (6mL /100m ²)	85mL	
Buffalo grass		WA only	1.2 (12mL /100m ²)	170mL	
Lucerne, Clover, Medics etc. grown for forage, hay or seed production	Annual Ryegrass, Winter grass, Barley grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue and Rats Tail Fescue	NSW, ACT, SA, WA, Tas only	1-1.5	SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July. WEEDS: Apply when weeds are at post or pre-emergent stages. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4 cm of soil. This is similar to 25mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatments is recommended.	
	Wild Oats, Perennial Ryegrass, Bentgrass, English Couch, Yorkshire Fog Grass, Chickweed, seedling Sorrel, Blackberry Nightshade, Patterson's Curse, seedling Wireweed		1.5-2		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

MIXING

Prior to pouring, shake container vigorously, then add the required quantity of EzyCrop Propyzamide 500 SC Herbicide to water in the spray vat while stirring or with agitators in motion. **DO NOT** mix with other pesticides.

RESISTANT WEEDS WARNING**GROUP****D****HERBICIDE**

EzyCrop Propyzamide 500 SC Herbicide is a member of the amide group of herbicides with diverse sites of action. For weed resistance management EzyCrop Propyzamide 500 SC Herbicide is a Group D herbicide. Some naturally occurring weed biotypes resistant to EzyCrop Propyzamide 500 SC Herbicide and other amide herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by EzyCrop Propyzamide 500 SC Herbicide or other amide herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, EzyCrop Pty Ltd accepts no liability for any losses that may result from failure of EzyCrop Propyzamide 500 SC Herbicide to control resistant weeds. Advice as to strategies and alternative treatments that can be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or EzyCrop Pty Ltd representative.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT
DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

PROTECTION OF LIVESTOCK
DO NOT graze sheep on treated areas or feed to poultry or pigs.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray drift onto nearby susceptible plants/crops, cropping land or pastures.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep from contact with fertilisers, insecticides, fungicides and seeds. Protect from frost.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site.

If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in accordance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat and elbow-length PVC gloves. After use, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre, Telephone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

The use of EzyCrop Propyzamide 500 SC Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by EzyCrop Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and EzyCrop Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

APVMA Approval No. 68069/57426

In a Transport Emergency Dial 000 Police or Fire Brigade

Batch Number:

Date of Manufacture:

