

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# EzyCrop Oxyfluorfen 240 Herbicide



ACTIVE CONSTITUENT: 240 g/L OXYFLUORFEN  
SOLVENTS: 606 g/L LIQUID HYDROCARBON  
108 g/L N-METHYL PYRROLIDONE

GROUP	<b>G</b>	HERBICIDE
-------	----------	-----------

For the selective control of certain broadleaf and grass weeds as  
per Directions for Use table

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE**

**CONTENTS: 10L, (20L)**

EzyCrop Pty Ltd  
2/22 Horne Street  
Elsternwick  
VIC 3185  
Australia

Tel: 03 9505 0044  
ACN: 156 476 827

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well-ventilated area. Store above 5°C.

Do not store for prolonged periods in direct sunlight.

DO NOT store near food, feedstuffs, fertilizers and seeds.

Rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SMALL SPILL MANAGEMENT**

Wear protective equipment (see SAFETY DIRECTIONS section). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb, as above, the wash liquid for disposal.

**SAFETY DIRECTIONS**

Will damage the eyes and irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length butyl rubber gloves and goggles. If product on skin, immediately wash area with soap and water. If product in eyes wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water.

**MATERIAL SAFETY DATA SHEET**


Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

**CONDITIONS OF SALE**

The use of EzyCrop Oxyfluorfen Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by EzyCrop Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and EzyCrop Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

APVMA Approval No.: 68131/57581

<b>UN NO. 3082</b>	ENVIRONMENTALLY HAZARDOUS SUBSTANCES LIQUID, N.O.S. (Contains oxyfluorfen) marine pollutant.	
In an Emergency <b>Dial 000</b> Police or Fire Brigade	PG III	<b>HAZCHEM 2Z</b>
	Specialist Poisons Advice: 13 11 26	



Batch No:

Date of Manufacture:

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **EzyCrop Oxyfluorfen 240 Herbicide**

**ACTIVE CONSTITUENT: 240 g/L OXYFLUORFEN**  
**SOLVENTS: 606 g/L LIQUID HYDROCARBON**  
**108 g/L N-METHYL PYRROLIDONE**

<b>GROUP</b>	<b>G</b>	<b>HERBICIDE</b>
--------------	----------	------------------

**For the selective control of certain broadleaf and grass weeds as  
per Directions for Use table**

**IMPORTANT: READ THIS LEAFLET BEFORE USE**

EzyCrop Pty Ltd  
2/22 Horne Street  
Elsternwick  
VIC 3185  
Australia

Tel: 03 9505 0044  
ACN: 156 476 827

**DIRECTIONS FOR USE**

EzyCrop Oxyfluorfen Herbicide can be used on weed-free soil to prevent germination of a wide variety of weeds or it can be applied to existing weeds at seedling stage especially with a tank mix partner to increase the variety of weeds controlled and/or the length of residual control. EzyCrop Oxyfluorfen Herbicide can also be added at a low rate as a 'spike' to glyphosate or paraquat and diquat/paraquat herbicides to improve knockdown.

**1. EzyCrop Oxyfluorfen Herbicide applied as a 'spike' with glyphosate OR with either paraquat or a diquat/paraquat mixture.**

**RESTRAINT:** DO NOT disturb weeds by cultivation or sowing for 1 day following application to annual weeds and 7 days for perennial weeds to ensure herbicide absorption, unless specified in the **CRITICAL COMMENTS**.

SITUATION	FOR WEEDS CONTROLLED & TIME OF APPLICATION	RATE of EzyCrop Oxyfluorfen Herbicide	CRITICAL COMMENTS
Summer and Winter Fallow – Before sowing	Refer to label of the glyphosate product (such as EzyCrop Glyphosate)	75 mL/ha plus a glyphosate product at its recommended label	Addition of EzyCrop Oxyfluorfen Herbicide to glyphosate products will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with glyphosate applied alone) and give control of annual nettles, ( <i>Urtica</i> spp.), barley grass, Patterson's curse, small-flowered mallow and storksbill. For rates of glyphosate, refer to the appropriate label. Read and follow all label directions. See SAFE SOWING INTERVALS section of this label.
Fruit & nut trees, vines including: Grapevines, Olive trees, Pome fruit (eg. apple, pear, nashi, quince), Stone fruit (eg. apricot, cherry, nectarine, peach, plum) Tree nuts (eg. almond, macadamia, pecan, walnut)	Refer to label of the glyphosate product (such as EzyCrop Glyphosate)	75 mL/ha plus a glyphosate product at its recommended label rate	Addition of EzyCrop Oxyfluorfen Herbicide to glyphosate products will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with glyphosate applied alone) and give control of annual nettles, ( <i>Urtica</i> spp.), barley grass, Patterson's curse, small-flowered mallow and storksbill. For rates of glyphosate, refer to the appropriate label. Read and follow all label directions. <b>DO NOT</b> apply the tank mix of glyphosate and EzyCrop Oxyfluorfen near trees or vines less than 3 years old unless they are effectively shielded from spray and spray drift.
	Refer to label of the paraquat or diquat/paraquat products (such as EzyCrop Paraquat or EzyCrop Paraquat/ diquat)	250 mL/ha plus a paraquat or diquat/paraquat product at its recommended label rate	Addition of EzyCrop Oxyfluorfen Herbicide in a tank mix with a paraquat or diquat/paraquat product will improve control of small flowered mallow, evening primrose and other weeds sensitive to EzyCrop Oxyfluorfen Herbicide. For the rate of the paraquat or diquat/paraquat product, refer to the appropriate label. Read and follow all label directions.

**2. EzyCrop Oxyfluorfen Herbicide applied to weed-free soil or weeds at seedling stage.**

**RESTRAINT:** If applying to weed seedlings, DO NOT disturb weeds by cultivation or sowing for 1 day following application to annual weeds and 7 days for perennial weeds to ensure herbicide absorption, unless specified in the CRITICAL COMMENTS.

CROP	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
<b>Brassica Crops</b> Broccoli, Cabbages, Cauliflower	Refer to Weeds Controlled list	Weed free soil (prior to crop transplanting)	1.5 to 2 L/ha	Apply EzyCrop Oxyfluorfen to prepared ground 4 to 7 days prior to transplanting. If soil is dry, irrigation or rainfall is required prior to transplanting for activation of EzyCrop Oxyfluorfen. Utilise transplanting techniques, which cause minimal soil disturbance.  Excessive soil disturbance will lessen herbicide activity. Use the higher rate in situations where weed pressure is known to be heavy.
		Weeds at 2 to 4 true leaf stage		When seedlings are present, apply as a tank mix with paraquat to give both knockdown and residual control.  A non-ionic surfactant should be used in the spray mixture at its recommended rate. Apply as a directed spray to avoid contact with coffee plants. Mature, established weeds must be eliminated by mechanical or chemical means prior to application.
<b>Coffee</b> (after transplanting or under established bushes)	Refer to Weeds Controlled list	Weed free soil	2 or 4 L/ha	Use the higher rate where longer residual activity (up to 4 months) is required.
		Weeds at 2 to 4 true leaf stage		When seedlings are present, apply as a tank mix with paraquat to give both knockdown and residual control.  A non-ionic surfactant should be used in the spray mixture at its recommended rate. Apply as a directed spray to avoid contact with coffee plants. Mature, established weeds must be eliminated by mechanical or chemical means prior to application.
<b>Duboisia</b> (after transplanting of young seedlings or after harvest of mature plants).	Refer to Weeds Controlled list (best results are achieved when applied to moist soil free of weeds)	Weed free soil	4 or 8 L/ha	Use the higher rate where longer residual control (up to 6 months) is required.  An 'Over-the-top' application will be tolerated.
		Weeds at the 4 to 6 true leaf stage		Recently germinated small seedling grasses and broadleaf weeds (4 to 6 true leaves) will be controlled with these rates.  Established larger weeds must be eliminated by mechanical or chemical means prior to the application of EzyCrop Oxyfluorfen.  Add a non-ionic surfactant at its recommended rate to enhance activity.
<b>Forestry Plantations:</b> <i>Eucalyptus</i> spp. and <i>Pinus</i> spp. (either before or preferably within 4 weeks of transplanting)	Refer to Weeds Controlled list	Weed free soil (either before or preferably within 4 weeks of transplanting)	3 or 4 L/ha	Under weed-free conditions, apply as a directed or 'over-the-top' spray. Disturbance of the herbicidal barrier on the soil surface at transplantation may reduce the length of weed control. If weed seedlings are present, apply as an 'over-the-top' spray.  In either situation, use the higher rate for longer residual control. EzyCrop Oxyfluorfen Herbicide can be applied in a tank mix with simazine to extend the spectrum and length of weed control.  <b>DO NOT</b> use this tank mix in Eucalyptus plantations grown on sands, with no clay or organic matter. The likelihood of foliar damage to trees (especially eucalypts) will increase if applied to foliage that has not hardened off and/or if the temperature exceeds 20°C. However EzyCrop Oxyfluorfen Herbicide is generally regarded as safe to commonly planted forestry species but the sensitivity of less common species should be tested on small areas before a large-scale application is made.
		Weeds at the 4 to 6 true leaf stage		Under weed-free conditions, apply as a directed or 'over-the-top' spray. Disturbance of the herbicidal barrier on the soil surface at transplantation may reduce the length of weed control. If weed seedlings are present, apply as an 'over-the-top' spray.  In either situation, use the higher rate for longer residual control. EzyCrop Oxyfluorfen Herbicide can be applied in a tank mix with simazine to extend the spectrum and length of weed control.  <b>DO NOT</b> use this tank mix in Eucalyptus plantations grown on sands, with no clay or organic matter. The likelihood of foliar damage to trees (especially eucalypts) will increase if applied to foliage that has not hardened off and/or if the temperature exceeds 20°C. However EzyCrop Oxyfluorfen Herbicide is generally regarded as safe to commonly planted forestry species but the sensitivity of less common species should be tested on small areas before a large-scale application is made.

Forestry Trees	Broadleaf weeds and grasses	Weeds at the 4 to 6 true leaf stage	4 L/ha Or 4mL/10m <sup>2</sup>	For the establishment of trees for approved farm practices such as wind breaks, erosion control, woodlots and forestry plantings. When applying as a post-plant spray, ensure spray is directed to the base of seedlings, or that seedlings are protected. <b>Do not apply</b> under hot or windy conditions.
Pyrethrum - as bare rooted transplants or seedlings	Refer to Weeds Controlled list (except chickweed)	Pre-plant incorporated into weed free soil worked to a fine tilth	4 or 6 L/ha	Apply prior to final soil preparation. The preferred implements for final soil preparation would be either a multiple tyne cultivator or rotary harrows. Use the 6 L/ha rate for heavy black clay soils only (as found in the Derwent and Coal River Valleys, Tasmania). EzyCrop Oxyfluorfen Herbicide will not provide consistent control of chickweed.
Pyrethrum - more than 4 leaves	Blackberry nightshade, Cleavers, Field bindweed, Fumitory, Groundsel, Sorrel, Volunteer potato, Wireweed	Emerged weeds present	100 to 150 mL/ha	
Pyrethrum - established crops, > 1 year old	As above plus Sow Thistle, Spear Thistle		200 mL to 4 L/ha	Apply when pyrethrum is > 10 cm rosettes. Apply rates of more than 1.0 L/ha ONLY between 1st of February and 31st of March. <b>DO NOT</b> apply later than 25 weeks before harvest.
Tobacco	Refer to Weeds Controlled list	Weed free soil	4 L/ha	Use to control weeds along spray lines only. <b>DO NOT</b> apply to tobacco crop. Apply to soil after solid-set irrigation system has been laid out in the field. EzyCrop Oxyfluorfen should be applied to moist soil. Where very small weeds (2 to 3 leaf) emerge prior to spraying, the addition of a non-ionic surfactant to the spray mixture is necessary for effective control. Should the weeds be more advanced, the addition of 2 L/ha diquat (200g/L) is required. Avoid spray drift.
Trees (Fruit & Nuts) and Vines at least 3 years old as a dormant application, including: Grapevines, Olive trees, Pome fruit (eg. apple, pear, nashi, quince), Stone fruit (eg. apricot, cherry, nectarine, peach, plum), Tree nuts (eg. almond, macadamia, pecan, walnut)	Refer to Weeds Controlled list	Weed free soil	3 or 4 L/ha	<b>DO NOT</b> apply EzyCrop Oxyfluorfen once bud swell has occurred. Apply to freshly cultivated, weed free soil. Use higher rate for longer residual control (up to 4 months). Where grass weeds are expected to be the major problem, or when control of a wider weed spectrum is needed, mix the lower rate with 4.5L/ha of an oryzalin (500g/L) product or 4.5 kg/ha of a napropamide (500 g/kg) product. Refer to product labels for crops, rates, states and weeds controlled and follow all label directions.

		Emerged weeds present (4-6 leaf stage)		<p><b>DO NOT</b> apply EzyCrop Oxyfluorfen once bud swell has occurred. Use the higher rate when longer residual control is required (up to 4 months). When young seedling grasses and/or broadleaved weeds are present, apply as a tank mix with glyphosate or paraquat or paraquat/diquat to obtain both knockdown and residual control. A non-ionic surfactant should be used at 0.1%v/v. Read and follow all label directions.</p> <p>Where weed growth is large and dense, weeds must be eliminated prior to application of EzyCrop Oxyfluorfen, using mechanical or chemical means.</p> <p><b>Macadamias:</b> Apply in 250 to 500 L water/ha. Apply after harvest to prevent spray contacting nuts. Avoid spray contact with the foliage and stem. <b>DO NOT</b> apply to nuts on the ground.</p>
Tropical and sub-tropical fruit crops (inedible peel), including; Avocado, Cherimoya, Custard apple, Durian, Feijoa, Guava, Jackfruit, Kiwi fruit, Longan, Lychee, Mango, Mangosteen, Papaya, Passionfruit, Persimmon, Rambutan, Star apple	Refer to Weeds Controlled List	Weed free soil	4 L/ha	Best results are obtained when applied to moist weed-free soil and followed by rainfall or irrigation.
		Emerged weeds present		<p>If weeds are present, EzyCrop Oxyfluorfen Herbicide should be applied as a tank mix with recommended rates of glyphosate or paraquat or diquat/paraquat.</p> <p>Read and follow all label directions.</p>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS:**

**GRAZING – DO NOT ALLOW LIVESTOCK TO GRAZE TREATED WEEDS**  
**HARVEST – NOT REQUIRED WHEN USED AS DIRECTED**

## WEEDS CONTROLLED BEFORE GERMINATION

Amsinckia ( <i>Amsinckia</i> spp.)	Prickly lettuce ( <i>Lactuca</i> spp.)
Barley grass ( <i>Hordeum leporinum</i> )	Red natal grass ( <i>Rhynchelytrum repens</i> )
Barnyard grass ( <i>Echinochloa</i> spp.)	Redshank ( <i>Amaranthus cruentus</i> )
Blackberry nightshade ( <i>Solanum nigrum</i> )	Ryegrass ( <i>Lolium</i> spp.)
Bladder Ketmia ( <i>Hibiscus trionum</i> )	Sesbania pea ( <i>Sesbania cannabina</i> )
Burrgrass ( <i>Cenchrus australis</i> )	Shepherd's purse ( <i>Capsella bursa-pastoris</i> )
Caltrop ( <i>Tribulus terrestris</i> )	Smallflower mallow ( <i>Malva parviflora</i> )
Capeweed ( <i>Arctotheca calendula</i> )	Soursob ( <i>Oxalis pes-caprae</i> )
Chickweed ( <i>Stellaria media</i> )	Sowthistle ( <i>Sonchus oleraceus</i> )
Crowsfoot grass ( <i>Eleusine indica</i> )	Starburr ( <i>Acanthospermum hispidum</i> )
Deadnettle ( <i>Lamium amplexicaule</i> )	Stinkgrass ( <i>Eragrostis cilianensis</i> )
Fat hen ( <i>Chenopodium album</i> )	Summer grass ( <i>Digitaria</i> spp.)
Giant pigweed ( <i>Trianthema portulacastrum</i> )	Thornapple ( <i>Datura stramonium</i> )
Liverseed grass ( <i>Urochloa panicoides</i> )	White eye ( <i>Richardia brasiliensis</i> )
Lovegrass ( <i>Eragrostis</i> spp.)	Wild mustard ( <i>Sisymbrium</i> spp.)
Pigeon grass ( <i>Setaria</i> spp.)	Wild radish ( <i>Raphanus raphanistrum</i> )
Pigweed ( <i>Portulaca oleracea</i> )	Wireweed ( <i>Polygonum aviculare</i> )

## SEEDLINGS

Amsinckia ( <i>Amsinckia</i> spp.)	Pigweed ( <i>Portulaca oleracea</i> )
Bellvine ( <i>Ipomoea</i> spp.)	Potato weed ( <i>Galinsoga parviflora</i> )
Capeweed ( <i>Arctotheca calendula</i> )	Redshank ( <i>Amaranthus cruentus</i> )
Common cotula ( <i>Cotula australis</i> )	Shepherd's purse ( <i>Capsella bursa-pastoris</i> )
Crowsfoot grass ( <i>Eleusine indica</i> )	Sowthistle ( <i>Sonchus oleraceus</i> )
Deadnettle ( <i>Lamium amplexicaule</i> )	Stinging nettle ( <i>Urtica urens</i> )
Groundsel ( <i>Senecio vulgaris</i> )	Stinkgrass ( <i>Eragrostis cilianensis</i> )
Liverseed grass ( <i>Urochloa panicoides</i> )	Wild radish ( <i>Raphanus raphanistrum</i> )

## GENERAL INSTRUCTIONS

EzyCrop Oxyfluorfen Herbicide is a selective herbicide for the control of certain annual grasses and broadleaf weeds in dormant apples, grapes, peaches, pears, plums, apricots, almonds, macadamia, Duboisia, tobacco, coffee, pyrethrum, winter cereals, tropical/sub-tropical crops, brassicas, onions, *Eucalyptus* and *Pinus* spp and other forestry trees applied either to a weed-free soil or to seedling weeds at the 4 to 6 true leaf stage.

EzyCrop Oxyfluorfen Herbicide applied to well prepared, weed free soil should not be disturbed or incorporated after application. Weed control for up to 6 months can be expected with high label rates, but spot treatment of escape weeds or perennial grasses may be necessary with knockdown herbicides.

When EzyCrop Oxyfluorfen Herbicide is applied to seedling weeds at the 4 to 6 leaf stage, a non-ionic surfactant such as BS-1000 should be added at recommended rates to improve activity, where suggested in the CRITICAL COMMENTS.

EzyCrop Oxyfluorfen Herbicide can also be used at low rates as a 'spike' to improve the weed spectrum of knockdown herbicides such as glyphosate and paraquat or diquat/paraquat mixtures.

## RESISTANT WEEDS WARNING

GROUP	G	HERBICIDE
-------	---	-----------

EzyCrop Oxyfluorfen Herbicide is a member of the diphenyl ether group of herbicides. The mode of action of EzyCrop Oxyfluorfen is to inhibit protoporphyrinogen oxidase. For weed resistance management EzyCrop Oxyfluorfen is a Group G herbicide.

Some naturally occurring weed biotypes resistant to EzyCrop Oxyfluorfen Herbicide and other Group G herbicides may exist through normal genetic variability in any weed population. The resistant



individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by EzyCrop Oxyfluorfen or other Group G herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, EzyCrop Pty Ltd accepts no liability for any losses that may result from the failure of EzyCrop Oxyfluorfen to control resistant weeds.

## **TIMING**

### **Residual Control**

For optimum residual weed control, EzyCrop Oxyfluorfen Herbicide should be applied to the soil surface prior to weed emergence after all other agricultural operations, such as mechanical cultivation and reshaping of irrigation furrows, have been completed. The area should be left undisturbed during the period of desired weed control. When applied to seedling weeds, they should be young and actively growing. Weed control for up to 6 months is expected but spot treatment, with knockdown herbicides, for escape weeds and perennial grasses may be necessary.

### **Post-emergence weed control**

For optimum post-emergence weed control, EzyCrop Oxyfluorfen herbicide + glyphosate tank mixes should be applied to small seedling weeds up to 4 - 6 true-leaf stage. Use of a non-ionic surfactant such as BS-1000 is recommended to improve activity. Weeds should be actively growing and free from environmental stress (drought, cold, insect attack, nutrient deficiency). Cultivation after treatment and prior to or at planting is beneficial for final fallow weed control.

## **MIXING**

Shake well before use. When using EzyCrop Oxyfluorfen alone, fill the spray tank, at least one-third full with clean water, add the recommended amount of EzyCrop Oxyfluorfen while the pump and agitator are running, then complete filling the spray tank.

A non-ionic surfactant, if required by label directions, should be added near the end of the filling process to minimise foaming.

If tank mixing with oryzalin (500g/L) or napropamide, add the product to a one-third filled tank, then the EzyCrop Oxyfluorfen during the filling operation. Maintain agitation during mixing and until spraying is completed.

When tank mixing with glyphosate formulations, paraquat, paraquat/diquat or diquat, add these after EzyCrop Oxyfluorfen during the filling operation.

### **EzyCrop Oxyfluorfen Herbicide + Glyphosate tankmixes**

Ensure thorough agitation when mixing, filling the spray tank and during application, irrespective of glyphosate formulation used. Follow recommended order and directions for tank mixing EzyCrop Oxyfluorfen and glyphosate. Use all spray mix immediately after preparation.

- DO NOT tank mix EzyCrop Oxyfluorfen Herbicide and glyphosate without agitation.
- DO NOT allow mix to stand unagitated.
- DO NOT store EzyCrop Oxyfluorfen Herbicide and glyphosate tank mixes.
- DO NOT mix other agrochemical products with EzyCrop Oxyfluorfen and glyphosate tank mixtures.

## **APPLICATION**

Spray equipment should be calibrated carefully before use.

EzyCrop Oxyfluorfen Herbicide should be applied uniformly with an accurately calibrated, low pressure herbicide sprayer, as a directed treatment to the base of tree and vine crops using flat fan or hollow cone nozzles. Complete coverage of seedling weeds is required for maximum knockdown effect. Ensure both weed foliage and the soil surface are sprayed.

Apply using a vehicle mounted boom, calibrate to deliver a droplet spectrum classification defined as medium by the ASABE Standard S572, using water volume of 250 to 500 litres per hectare for bare soil or 100 to 1350 litres per hectare when seedling weeds (4 to 6 leaf stage) are treated. Use the higher volumes where weed density is high.

Tank mixtures of 75 mL/ha of EzyCrop Oxyfluorfen Herbicide with glyphosate herbicides should be applied in 30 to 200 litres spray volume per hectare. For maximum residual control, EzyCrop Oxyfluorfen Herbicide should NOT be incorporated or disturbed after application.

## **CROP SAFETY**

EzyCrop Oxyfluorfen may be applied as directed and/or shielded spray around dormant peach, plum, apricot, almond, apple and pear trees and grape vines of all ages when applied at rates of less than 1.0 L/ha.

When applied at 3.0 L/ha and above, the trees and grape vines should be at least 3 years of age. Do NOT apply EzyCrop Oxyfluorfen once bud swell has occurred when using rates greater than 1.0 L/ha. Duboisia seedlings and mature plants will tolerate 'over-the-top' applications of EzyCrop Oxyfluorfen.

### SAFE SOWING INTERVALS

EzyCrop Oxyfluorfen Herbicide at up to 75 mL/ha may be safely applied 1 day prior to planting broadacre crops such as cereals (wheat, barley, oats, triticale), canola, pulses (lupins, faba beans, field peas) and undersown pastures (lucerne, clover, medics, ryegrass, phalaris, cocksfoot) and 7 days minimum prior to planting cotton or soybeans, provided minimum tillage planting equipment is used with minimal soil disturbance. Inversion, mixing of surface soil with that in the planting zone or covering seed with treated soil may result in injury to emerging crop seedlings.

**Avoid covering the seed with soil treated with EzyCrop Oxyfluorfen Herbicide during the planting operation to minimise crop injury.**

EzyCrop Oxyfluorfen Herbicide has residual soil activity, especially when applied at rates greater than 75 mL/ha and on small-seeded horticultural crops. Plant back intervals in the following table for horticultural crops must be observed if more than 75 mL/ha EzyCrop Oxyfluorfen Herbicide has been applied.

### Safe Sowing Intervals for Horticultural Crops (days)

EzyCrop Oxyfluorfen Herbicide rate	Up to 75 mL/ha	1 L/ha
Beans	7	60
Brassicas	14	90
Capsicums	14	90
Carrots	14	90
Cucurbits	14	60
Lettuce	14	90
Potatoes	7	60
Tomatoes	14	60

### COMPATIBILITY

EzyCrop Oxyfluorfen is compatible with glyphosate products (with agitation), oryzalin (500g/L) paraquat, paraquat/diquat, diquat and glufosinate products.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

### PROTECTION OF LIVESTOCK

Use with care when applying in areas frequented by stock.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

This product is highly toxic to wildlife and fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Use care when applying in areas frequented by wildlife or adjacent to any body of water. DO NOT apply when weather conditions favour drift from target areas.

### STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Store above 5°C. Do not store for prolonged periods in direct sunlight. DO NOT store near food, feedstuffs, fertilizers and seeds. Rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Will damage eyes and will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length butyl rubber gloves and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hand after use. After each day's use, wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

**CONDITIONS OF SALE**

The use of EzyCrop Oxyfluorfen Herbicide being beyond the control of the manufacturer; no warranty expressed or implied is given by EzyCrop Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and EzyCrop Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

APVMA Approval No: 68131/57581