

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



EzyCrop Bromoxynil 200 Herbicide

**ACTIVE CONSTITUENT: 200 g/L BROMOXYNIL present as
the n-octanoyl ester**

SOLVENT: 616 g/L LIQUID HYDROCARBON

GROUP

C

HERBICIDE

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE OPENING OR USING
THIS PRODUCT**

**CONTENTS: 20 LITRES
(5L, 200L & 1000L)**

EzyCrop Pty Ltd
2/22 Horne Street
Elsternwick
VIC 3185
Australia

Tel: 03 9505 0044
ACN: 156 476 827

GENERAL INSTRUCTIONS

Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage and Disposal, Safety Directions and First Aid instructions. Use only at recommended growth stages of crops and weeds. DO NOT use on crops other than recommended.

STORAGE AND DISPOSAL**(5L, 20L and 200L only)**

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple or pressure rinse container for disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. Deliver empty packaging for appropriate disposal to an approved waste management facility. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots.

Empty containers and product should not be burnt.

For refillable containers (1000L only):

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight.

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow length PVC gloves and face shield or goggles. If product is in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet that can be obtained from the supplier.

CONDITIONS OF SALE

The use of EzyCrop Bromoxynil 200 Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by EzyCrop Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and EzyCrop Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

APVMA Approval No: 68188 / 57736

In a Transport Emergency Dial 000 Police or Fire Brigade

Batch No:
DOM

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

EzyCrop Bromoxynil 200 Herbicide

**ACTIVE CONSTITUENT: 200 g/L BROMOXYNIL present as
the n-octanoyl ester**
SOLVENT: 616 g/L LIQUID HYDROCARBON

GROUP

C

HERBICIDE

**For the control of weeds in wheat, oats, barley, cereal rye and
triticale (alone or undersown), linseed, clover and lucerne pastures,
turf, fallow, non-crop, roadsides and right-of-way as specified in
the Directions for Use table.**

READ THIS LEAFLET BEFORE OPENING OR USING THIS PRODUCT

APVMA Approval No: 68188 / 57736

EzyCrop Pty Ltd
2/22 Horne Street
Elsternwick
VIC 3185
Australia

Tel: 03 9505 0044
ACN: 156 476 827

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT apply by mister machines.

DO NOT apply if crops or weeds are stressed due to excessively dry or moist conditions.

DO NOT apply to diseased or frost-affected crops or if frosts are imminent.

DO NOT apply if rain is expected within 3 hours.

DO NOT apply to lucerne crops or clover-based pastures or turf in temperatures above 20°C or when temperatures above 20°C may follow for some days after application.

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate /ha	Critical Comments
Wheat, Barley, Cereal Rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medic (Vic only); Linseed; Clover or Lucerne Pastures; Lotus Seed or Forage Crops (Qld only)	Amsinckia (<i>Amsinckia spp.</i>), Capeweed (<i>Arctotheca calendula</i>) Black Bindweed (Climbing Buckwheat) (<i>Fallopia convolvulus</i>) Chamomile (<i>Matricaria matricarioides</i>), Common Peppergrass (<i>Lepidium hyssopifolium</i>), Corn Gromwell (Ironweed, Sheepweed), (<i>Buglossoides arvensis</i>) Fat Hen (<i>Chenopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mountain Sorrel (<i>Oxalis acetosella</i>) Purple Calandrinia (<i>Calandrinia menziesii</i>), Saffron Thistle (<i>Carthamus lanatus</i>), Shepherd's Purse (<i>Capsella bursa-pastoris</i>), Tree Hogweed (<i>Polygonum patulum</i>), Variegated Thistle (<i>Silybum marianum</i>)	Up to the 4 leaf stage or when plants are no more than 35 mm in diameter	All states	1.4 L	CROP STAGE: Wheat, Barley, Cereal, Rye, Oats and Triticale: 3 leaf to fully tillered (Zadoks Scale Z13-30). Linseed: 50 to 150 mm high. Lucerne and Clover: After seedlings have one trifoliate leaf or before there is 150 mm regrowth in established stands. Lotus: Apply post-emergence at least 4 weeks after sowing (crop seedlings with a min. of 2-6 true leaves). APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weeds is essential. Apply only in the Autumn and Winter (mid-April to end of August). For optimum control, apply only when the weeds are actively growing and before the weeds are shielded by the crop. CAUTION: Lucerne and Clover: This product will cause slight leaf burning of undersown lucerne and Clover. DO NOT apply to Persian Clover or Berseem Clover. Linseed: DO NOT spray linseed by aerial application. Lotus: Some initial scorching of leaves may occur in some cases. Medics: Application to some cultivars of Medics may result in unacceptable crop damage in some situations. # Effective control of Paterson's Curse and Fumitories will only be obtained under good growing conditions. ## Where Wireweed is growing under stress, it has been found to be less susceptible.
	Mexican Poppy (<i>Argemone ochroleuca</i>), Three-cornered Jack (Doublegee, Spiny Emex), (<i>Emex australis</i>) Turnip Weed (Giant Mustard), (<i>Rapistrum rugosum</i>)	Up to the 6 leaf stage or when plants are no more than 50 mm in diameter		2.0 L	

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate /ha	Critical Comments
As above	Wild Radish (<i>Raphanus raphanistrum</i>), Wild Mustard (<i>Sisymbrium spp.</i>), Wild Turnip (<i>Brassica tournefortii</i>), Wireweed## (<i>Polygonum aviculare</i>), Paterson's Curse# (<i>Salvation Jane</i>) (<i>Echium plantagineum</i>)	As above	As above	As above	As above
	Fumitories# (<i>Fumaria spp.</i>)		Qld, NSW, Vic, SA, WA, ACT only		
	Pheasant's Eye (<i>Adonis microcarpa</i>) Birds Eye (<i>Carrot Weed</i>) (<i>Cotula australis</i>)	Up to the 4 true leaf stage	SA only	700 mL	DO NOT use where undersown medics occur.
		Greater than the 4 true leaf stage.		1.4 L	
	Fireweed (<i>Senecio spp.</i>)	Seedling	Qld, NSW, ACT only	1.4 L	Apply during the Autumn-Winter period when weed are young and actively growing.
		Early flowering stage		2.8 L	
Wheat, Barley, Cereal Rye, Oats and Triticale: including those undersown with Clover, Lucerne or Medics; Clover Seed Crops	Amsinckia (<i>Amsinckia spp.</i>), Capeweed (<i>Arctotheca calendula</i>), Chamomile (<i>Matricaria matricarioides</i>), Charlock (<i>Sinapis arvensis</i>), Corn Gromwell (Ironweed, Sheepweed), (<i>Buglossoides arvensis</i>) Fat Hen (<i>Chenopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Hexham Scent (King Island Melilot) (<i>Melilotus indicus</i>), Horned Poppy (<i>Glaucium flavum</i>), Indian Hedge Mustard (<i>Sisymbrium</i>)	Up to the 5 leaf stage or when weed plants are no more than 50 mm in diameter	WA only	750 mL/ha plus 440 mL of Amicide* 625/ha or plus 700 mL of MCPA 500/ha	CROP STAGE: CEREALS: 3 leaf to fully tillered. (Zadoks Scale Z13-30). CLOVER: Apply after clover seedlings have at least three trifoliate leaves. APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Apply only in the Autumn and Winter (mid-April to end of August). Apply only when the weeds are actively growing and before the weeds are shielded by the crop. CAUTION: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT apply to Persian clover or Berseem clover. A slight burning of clover may occur. DO NOT use 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or Medics unless some damage is acceptable. (continued on next page)
			Vic, SA only	1.4 L /ha plus 440 mL of Amicide* 625/ha or plus 700 mL of MCPA 500/ha	

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate /ha	Critical Comments
As above	<i>orientale</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mexican Poppy (<i>Argemone mexicana</i>), Mintweed (<i>Salvia reflexa</i>), Purple Calandrinia (<i>Calandrinia menziesii</i>) Paterson's Curse (Salvation Jane) (<i>Echium plantagineum</i>) Rough Poppy (<i>Papaver hybridum</i>), Saffron Thistle (<i>Carthamus lanatus</i>), Shepherd's Purse (<i>Capsella bursa-pastoris</i>), Slender Thistle (<i>Carduus tenuiflorus</i>), Turnip Weed (Giant Mustard) (<i>Rapistrum rugosum</i>), Wild Radish (<i>Raphanus raphanistrum</i>)	As above	As above	As above	This product may be tank mixed with Hoegrass*/Nugrass*.
	Common Peppergrass (<i>Lepidium hyssopifolium</i>), Tree Hogweed (<i>Polygonum patulum</i>)		Vic only		
	Black Bindweed (Climbing Buckwheat) (<i>Polygonum convolvulus</i>)		Up to the 8 leaf stage		
Wheat, Barley, Oats and Triticale (alone or undersown with Lucerne and Clover)	Common Sowthistle (<i>Sonchus oleraceus</i>)				APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application. Good coverage of the spray on the weed is essential. Apply only in the Autumn and Winter (mid-April to end of August). For optimal control, apply when the weeds are actively growing and before the weeds are shielded by the crop. CAUTION: Lucerne and Clover: This product will cause slight leaf burning of undersown Lucerne and Clover. DO NOT apply to Persian Clover and Berseem Clover. (continued below)

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate /ha	Critical Comments
		Up to 4 leaves	QLD, northern NSW	1.4 L/ha	Strictly for use on small weeds early in crop before shading out. CROP STAGE Wheat, Barley, Oats and Triticale: 3-leaf to mid-tillering (Z13-Z23). Use a minimum spray volume of 70L/ha. If Broup B resistance is suspected, or weeds have grown through a previous application of chlorsulfuron, apply 2.1L/ha.
		4-8 leaves		2.1 L/ha	CROP STAGE Wheat, Barley, Oats and Triticale: 3-leaf to full-tillering (Z13-Z30). This product will cause slight leaf burning of undersown Lucerne and Clover.
Wheat, Barley, Oats and Triticale	Three-cornered Jack (Doublegee, Spiny Emex) (<i>Emex australis</i>) Wireweed (<i>Polygonum Aviculare</i>) Variegated Thistle (<i>Silybum marianum</i>)	Up to the 4-leaf stage or when plants are no more than 50mm in diameter	Qld, NSW, Tas, SA, ACT only	1.4 L /ha plus 440 mL of Amicide* 625/ha or plus 700 mL of MCPA 500/ha	CROP STAGE: 5 leaf to fully tillered (Zadoks scale Z15-30); Apply under good growing conditions. BOOM SPRAYING: Use a minimum of 220 L spray per hectare on Black Bindweed when crop is greater than 300 mm high. CAUTION: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT mix with 2,4-D or MCPA tank mix if wheat, oats, barley or triticale are undersown with lucerne, clover or medics unless some damage is acceptable. DO NOT apply to medics (WA only). (continued on next page) (continued from above)
	Fumitories (<i>Fumaria spp.</i>)		Qld, NSW, ACT, Vic only		
	Wild Mustard (<i>Sisymbrium spp.</i>), Wild Turnip (<i>Brassica tournefortii</i>)		Qld, Vic only		
As above	Cleavers	2-4 stems + 1-3 whorls of leaves /stem	Qld, Vic, NSW, ACT, Tas only	As above	This product may be tank mixed with Hoegrass*/Nugrass or control of Annual Ryegrass and Wild Oats in Wheat and Barley only. DO NOT use 2,4-D or MCPA tank mix if using a tank with diclofop-methyl.
	Three-cornered Jack (Doublegee, Spiny Emex) (<i>Emex australis</i>) Wireweed (<i>Polygonium</i>)	4 – 5 leaf stage	WA only	2.1 L plus 550 mL of 500 g/L 2,4-D amine or plus 700 mL of 500 g/L MCPA amine	APPLICATION: Refer to General Instructions. Avoid application when maximum daily temperatures above 20°C occur, or are likely to occur for a few days after application.
	Fumitories (<i>Fumaria spp.</i>)		Qld, Vic, NSW, ACT, SA, Tas only		
			Qld, NSW, Vic, SA only		
	Black Bindweed (<i>Polygonum convolvulus</i>) (Climbing Buckwheat), Chamomile (<i>Matricaria matricarioides</i>),	5 to 8 leaf stage	All states		CROP STAGE: 5 leaf to fully tillered (Zadoks scale Z15-30); Apply under good growing conditions. BOOM SPRAYING: Use a minimum of 220 L spray per hectare on Black Bindweed when crop is greater than 300 mm high.

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate /ha	Critical Comments
	Fat Hen (<i>Chemopodium album</i>), Field Madder (<i>Sherardia arvensis</i>), Lesser Swinecress (<i>Coronopus didymus</i>), Mountain Sorrel (<i>Calandrinia menziesii</i>), Paterson's Curse (<i>Echium plantagineum</i>) (Salvation Jane), Shepherd's Purse (<i>Capsella bursa-pastoris</i>), Variegated Thistle (<i>Silybum marianum</i>)				CAUTION: Some crop damage and/or yield loss may occur in Olympic and Shortim wheat cultivars. DO NOT mix with 2,4-D or MCPA tank mix if wheat, oats, barley or tritcale are undersown with lucerne, clover or medics unless some damage is acceptable. DO NOT apply to medics (Western Australia only). This product may be tank mixed with diclofop-methyl for control of Annual Ryegrass and Wild Oats in Wheat and Barley only. DO NOT use 2,4-D or MCPA tank mix if using a tank with diclofop-methyl.
Fallow & Non-Crop	Cowvine or Peachvine (<i>Ipomea lonchophylla</i>)	Up to 6 leaves	Qld, NSW, ACT only	1.4 L	<u>For all fallow and non-crop situations</u> Complete coverage is essential, plants which are shaded by stubble or other weeds will not be controlled. Weeds that are drought stressed and sprayed under hot conditions may not be adequately controlled. The addition of glyphosate will provide broader spectrum weed control when needed. Refer to the compatibility section for appropriate glyphosate products. The rate of glyphosate used should be robust for the conditions and sufficient to control the hardest-to-kill weed in the situation. For general fallow weed control application volumes above 50L/ha are recommended.
		Up to 8 leaves		2.1 L	
	Climbing Buckwheat (<i>Fallopia convolvulus</i>)	Cotyledon to mature, vining plants		1.5 L/ha alone or 1 – 1.5 L/ha plus glyphosate	
	Volunteer conventional cotton (<i>Gossypium hirsutum</i>)	Cotyledon to 6 leaves	Qld, NSW, WA, NT only	1.5 L/ha alone or 1 – 1.5 L/ha plus glyphosate	
	Volunteer RoundUp Ready* cotton (<i>Gossypium hirsutum</i>)	Cotyledon to 6 leaves		1.5 L/ha alone or plus glyphosate	
Pastures	Adonis (Pheasants Eye), Birds Eye (Carrot Weed)	Up to 4 leaves	SA only	700mL	Apply in May-June. Medics may be severely scorched at the low rate and killed at the high rate. Do not apply to polymorpha medics. Avoid application to clover-based pastures in temperatures over 20°C or when temperatures above 20°C may follow for some days after application, as seedling mortality may occur and established plants may be damaged.
		More than 4 leaves		1.4L	
	Doublegee, Wild Radish	Up to 4 leaves	WA only	1.5L	
	Matricaria	Up to 3 leaves		750mL	

Situation & Crop	Weeds Controlled	Weed Stage	State	Rate /ha	Critical Comments
	Fireweed	Seedling	NSW, Qld, ACT only	1.4L	Apply with low volume boom spray during autumn/winter, when weeds are young and actively growing.
		Early flowering		2.8L	
Pastures, Roadsides, Rights-of-way	African Daisy (<i>Senecio pterophorus</i>)	Up to and including the flowering stage	SA only	300 mL per 100 L water	The plants, including the stems, must be thoroughly wet to the point of run-off. Apply when the plants are actively growing. A non-ionic wetter should be added at 125 mL per 100 litres of spray. A follow-up treatment may be necessary for large, well-established plants. Ensure an overall spray coverage on weed seedlings.
	Boneseed/Bitou Bush (<i>Chrysanthemoides monilifera</i>)	Seedling only	Vic, Tas only	160 mL per 100 L water	

Situation & Crop	Weeds Controlled	State	Rate			Critical Comments
Turf Established couch, (including Qld Blue), Buffalo, Bent, Paspalum, Kikuyu, Fescue and Rye lawns	Bindy-eye (<i>Calotis hispidula</i>),	All States	6L/500 L water per ha	High Volume	Knapsack	Apply evenly over the area to be treated. A slight browning of grasses, especially Paspalum, may occur in hot conditions.
	Jo-jo (Onehunga) (<i>Soliva pterosperma</i>), Cudweed (<i>Gnaphalium spp.</i>)			300 mL per 100 L	60 mL/100m ² per 15L	

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

WITHHOLDING PERIODS

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION

CROP HARVEST: NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS:

Before opening, carefully read Directions for Use, Precautionary and Protection Statements, Storage and Disposal, Safety Directions and First Aid Instructions. Use only at recommended growth stages of crops and weeds. **DO NOT** use on crops other than recommended.

MIXING INFORMATION

To ensure even mixing, half fill the spray tank with clean water and add the required amount of this product. Agitate thoroughly, then add the remainder of the water. Agitate again before spraying commences.

BOOM SPRAYER

Apply 50 to 200 litres of water per hectare (WA: 50 to 100L). Calibrate the machine before use to ensure that the correct application rate is obtained. Check height of boom above the crop to ensure that the spray is evenly distributed. A minimum of 160 litres and preferably 200 litres of water per hectare is recommended where the weed infestation is heavy or the crop cover is dense.

DO NOT use rain-drop or air-induction nozzles as coverage of weeds may not be adequate.

AERIAL APPLICATION

Apply a minimum of 22 litres of water per hectare. Markers should always move upwind before aircraft passed overhead. Application by Micronair equipment may give variable results.

EQUIPMENT MAINTENANCE

After use wash equipment thoroughly and rinse several times prior to re-use. Wash sprayer thoroughly with clean water after use. If using tank mixtures with 2,4-D amine or MCPA 250 apply with equipment that will not be used in 2,4-D and MCPA susceptible crops, particularly cotton and vines.

COMPATIBILITY

This product may be mixed in the spray tank with one of the following herbicides and insecticides in wheat and barley only:

metsulfuron, chlorsulfuron, diclofop-methyl, 2,4-D amine, MCPA 250, diflufenican, dicamba, chlorpyrifos and omethoate. Glyphosate products, including EzyCrop products.

As formulations of other manufacturer's products are beyond the control of EzyCrop Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

RESISTANT WEEDS WARNING

GROUP	C	HERBICIDE
-------	---	-----------

EzyCrop Bromoxynil 200 Herbicide is a member of the nitriles group of herbicides. The product has the inhibitors of photosynthesis at Photosystem II mode of action. For weed resistance management, EzyCrop Bromoxynil 200 Herbicide is a Group C herbicide.

Some naturally occurring weed biotypes resistant to EzyCrop Bromoxynil 200 Herbicide and other Group C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by EzyCrop Bromoxynil 200 Herbicide or other Group C herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, EzyCrop Pty Ltd accepts no liability for any losses that may result from the failure of EzyCrop Bromoxynil 200 Herbicide to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Drift Warning: DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Note: Maximum daily temperature at or above 20°C at application, or on days following treatment may result in seedling mortality and established plants may be damaged.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish.

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

(5L, 20L and 200L only)

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

Triple or pressure rinse container for disposal. Add rinsings to the spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. Deliver empty packaging for appropriate disposal to an approved waste management facility. If no landfill is available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots.

Empty containers and product should not be burnt.

For refillable containers (1000L only):

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Product is poisonous if inhaled or swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow length PVC gloves and face shield or goggles. If product is in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. (Phone Australia 13 11 26). If swallowed, **DO NOT** induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet that can be obtained from the supplier.

CONDITIONS OF SALE

The use of EzyCrop Bromoxynil 200 Herbicide being beyond the control of the manufacturer, no warranty expressed or implied is given by EzyCrop Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and EzyCrop Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

* Other registered trademarks.

APVMA Approval No: 68188 / 57736

**In a Transport Emergency
Dial 000
Police or Fire Brigade**