

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

EzyCrop Diflufenican + MCPA Selective Herbicide



ACTIVE CONSTITUENTS:

250 g/L MCPA (present as the ethyl hexyl ester)

25 g/L Diflufenican

SOLVENTS:

325 g/L LIQUID HYDROCARBONS

150 g/L N-methyl-2-pyrrolidone

GROUP	FI	HERBICIDE
-------	-----------	-----------

For the control of certain broadleaf weeds in winter cereals and clover as specified in the Directions for Use Table

IMPORTANT:

READ THE ATTACHED BOOKLET THOROUGHLY BEFORE
OPENING OR USING THE PRODUCT

Contents: 5L (20L, 100L, 1000L).

APVMA Approval No.: 68195/57750

EzyCrop Pty Ltd
2/22 Horne Street
Elsternwick
VIC 3185
Australia
Tel: 03 9505 0044
ACN: 156 476 827

DIRECTIONS FOR USE

For complete directions for use please read the attached leaflet.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area.
DO NOT store for prolonged periods in direct sunlight.

5L, 20L containers only:

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Re-fillable containers:

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured.

Empty product as required into application equipment. Do not attempt to breach the valve system or filling point, or contaminate the container with water or other products. Ensure that the equipment used in transfer of the product is disconnected, triple rinsed and drained after each use.

When the container is empty, close all caps and valves and return the container to the point of purchase.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, DO NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for EzyCrop Diflufenican + MCPA Selective Herbicide is available from EzyCrop Pty Ltd on request.

CONDITIONS OF SALE

EzyCrop Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on EzyCrop Pty Ltd's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of EzyCrop Pty Ltd has any authority to add to or alter these conditions.

* Other trademarks

BATCH NO:

DATE OF MANUFACTURE:

In an Emergency Dial 000 Police or Fire Brigade.

CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

EzyCrop Diflufenican + MCPA Selective Herbicide

ACTIVE CONSTITUENTS:

250 g/L MCPA (present as the ethyl hexyl ester)

25 g/L Diflufenican

SOLVENTS:

325 g/L LIQUID HYDROCARBONS

150 g/L N-methyl-2-pyrrolidone

GROUP	FI	HERBICIDE
--------------	-----------	------------------

For the control of certain broadleaf weeds in winter cereals and clover as specified in the Directions for Use Table

IMPORTANT:

**READ THE ATTACHED BOOKLET THOROUGHLY BEFORE
OPENING OR USING THE PRODUCT**

APVMA Approval No.: 68195/57750

EzyCrop Pty Ltd
2/22 Horne Street
Elsternwick
VIC 3185
Australia
Tel: 03 9505 0044
ACN: 156 476 827

DIRECTIONS FOR USE

Restraints:

DO NOT apply to crops or weeds which are stressed due to dry or excessively moist conditions.

DO NOT apply to crops under stress due to disease or insect damage.

DO NOT apply to frost-affected crops or if frosts are imminent.

DO NOT apply when heavy rain is expected within 4 hours.

CROP	WEEDS CONTROLLED	STAGE OF WEED GROWTH	STATE	RATE/ha	CRITICAL COMMENTS
<p>Cereals Wheat, barley, oats, triticale, cereal rye (including cereals undersown with clover)</p> <p>Pasture Newly sown and established clover-based pasture, clover for hay and seed production</p>	Wild Radish	Up to the 2 leaf stage and not more than 60mm in diameter	WA only	250mL	<p>CROP STAGE Cereals Up to 750mL (3 leaf to fully tillered stage - Z13 to 30). Over 750mL (5 leaf to fully tillered stage - Z15 to 30). Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing). WA only: DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15). Warning: EZYCROP Diflufenican + MCPA Selective Herbicide may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to "Crop Tolerance" section of General Instructions).</p> <p>Clover Application is recommended prior to the eighth trifoliate leaf stage; however, applications prior to the third leaf stage may result in crop damage especially under stressed conditions and in sandy soils. DO NOT apply to Annual Medics or lucerne. Warning: EZYCROP Diflufenican + MCPA Selective Herbicide may cause transient crop yellowing of clover, and may affect growth and seed set of some varieties of clover. (Refer to "Crop Tolerance" section of General Instructions).</p> <p>WEED STAGE Apply when weeds are actively growing. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as: * high crop and weed density; * late season germinations; * abnormal weed growth (including early flowering), higher rates of product (up to the maximum rate of application specified for that weed) may be required. EZYCROP Diflufenican + MCPA Selective Herbicide will not effectively control: * regrowth of suppressed weeds;</p>
		Up to the 4 leaf stage and not more than 120mm in diameter	ALL STATES	500mL	
		Up to the 6 leaf stage and not more than 150mm in diameter		750mL	
		Up to the 8 leaf stage and not more than 180mm in diameter		1.0L	
	Charlock, Hedge Mustard, Indian Hedge Mustard, Shepherd's Purse, Turnip Weed, Wild Turnip	Up to the 2 leaf stage and not more than 60mm in diameter	ALL STATES	500mL	
		Up to the 4 leaf stage and not more than 120mm in diameter		750mL	
		Up to the 6 leaf stage and not more than 150mm in diameter		1.0L	
	London Rocket	Up to the 5 leaf stage and not more than 120mm in diameter	QLD only	750mL	
	Ward's Weed		SA only		
	Capeweed	Up to the 2 leaf stage and not more than 60mm in diameter	ALL STATES	500mL	
		Up to the 4 leaf stage and not more than 120mm in diameter		1.0L	
	Crassula	Up to the 2 leaf stage	ALL STATES	500mL	
		Up to the 4 leaf stage		750mL	
	Prickly Lettuce	Up to the 2 leaf stage	ALL STATES	500mL	
		Up to the 4 leaf stage		750mL	
		Up to the 6 leaf stage		1.0L	
	Dense-flower Fumitory	Up to the 2 leaf stage	ALL STATES	750mL	
	Corn Gromwell, Saffron Thistle, Toad Rush		ALL STATES	1.0L	
	Deadnettle		NSW, VIC, SA only		
Sorrel	Up to the 2 leaf stage	VIC only	1.0L		
Canola (rape)	Up to the 4 leaf stage	ALL STATES	500mL		
Purple Goosefoot	Up to the 6 leaf stage	QLD only	500mL		
Turnip Weed, Wild Turnip	Cotyledon to 2 leaf stage	NSW only (West of Newell Hwy.) SA only (Eyre peninsula north of the line between Venus Bay and Cowell)	350mL		

<p>Cereals Wheat, barley, oats, triticale, cereal rye</p>	<p>Fumitory</p>	<p>2 - 6 leaf stage</p>	<p>ALL STATES</p>	<p>500mL + 200ml Terbutryn (500 g/L)</p>	<p>* transplanted weeds; * regrowth from rhizomes or roots; * weeds growing under stress from previous herbicide applications.</p>	
<p>GRAZING Efficacy on larger weeds will be improved by grazing with normal levels of stock after the 7 day withholding period. Refer to 'Protection of Livestock' for grazing precautions.</p>						
<p>SUPPRESSION OF THE FOLLOWING WEEDS</p>						
<p>Cereals Wheat, barley, oats, triticale, cereal rye (including cereals undersown with clover)</p>	<p>Saffron Thistle</p>	<p>Up to the 6 leaf stage</p>	<p>ALL STATES</p>	<p>1.0 L</p>	<p>APPLICATION Activity of this product will be reduced if weeds are stressed. Optimum results will be obtained if good soil moisture exists at and after application. Where crop or weed density is high, water volume should be increased. WILD RADISH EZYCROP Diflufenican + MCPA Selective Herbicide will provide residual control of Wild Radish for up to 4 weeks after application. Effective residual activity of this product may be reduced where: * rates lower than 1.0L/ha are used; * dry conditions prevail; * poor coverage of the soil surface is achieved; * crop is planted in non-wetting sand; * soils contain a high content of organic matter. Optimum results will be obtained if good soil moisture exists at and after application.</p>	
<p>Pasture Newly sown and established clover based pasture, clover for hay and seed production</p>	<p>Chickweed, Fireweed, Hexham Scent (King Island Melilot), Iceplant, Mouse-eared Chickweed, night-scented Stock, Paterson's Curse, Peppergrass, Skeleton Weed, Long Storksbill, Volunteer Lupins</p>	<p>Up to the 4 leaf stage</p>				<p>750 mL</p>
<p></p>	<p>Wireweed (Hogweed)</p>	<p>Up to the 2 leaf stage</p>				<p>1.0 L</p>
<p>Cereals Wheat, barley, oats, triticale, cereal rye</p>	<p>Wild Radish</p>	<p>Up to the 4 leaf stage and not more than 120mm in diameter</p>	<p>ALL STATES</p>	<p>350mL plus 200mL MCPA LVE (500g/L)</p>	<p>Refer also to all Critical Comments relating to weed stage, grazing, application and wild radish above. * Reduced efficacy (suppression only) may be achieved on Wild Radish larger than 8 leaf or greater than 180 mm in diameter.</p>	
<p></p>	<p></p>	<p>Up to the 6 leaf stage and not more than 150mm in diameter</p>		<p>500mL plus 200mL MCPA LVE (500g/L)</p>		

		Up to the 8 leaf stage and not more than 180mm in diameter*		500mL plus 400mL MCPA LVE (500g/L)	<p>DO NOT use this tank-mix if cereals are undersown with lucerne or annual medics.</p> <p>Crop Stage</p> <p>EZYCROP Diflufenican + MCPA Selective Herbicide 350mL + MCPA LVE 200m L: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30).</p> <p>EZYCROP Diflufenican + MCPA Selective Herbicide 500mL + MCPA LVE 200m L: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30).</p> <p>EZYCROP Diflufenican + MCPA Selective Herbicide 500mL + MCPA LVE 400mL: Apply from 5 leaf stage to fully tillered (Zadok's Z15 to Z30).</p> <p>Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing).</p> <p>WA only: DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15).</p> <p>Warning: EZYCROP Diflufenican + MCPA Selective Herbicide may cause transient crop yellowing of cereals. Some</p>
--	--	---	--	------------------------------------	---

varieties of oats have not been tested. (Refer to 'Crop Tolerance' section of General Instructions). Observe instructions also L.V.E. MCPA product label.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

CROP HARVEST:
ALL CROPS:

NOT REQUIRED WHEN USED AS DIRECTED.

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

WEED LIST

Common name	Scientific name
Canola (rape)	Brassica napus
Capeweed	Arctotheca calendula
Charlock	Sinapis arvensis
Chickweed	Stellaria media
Common Sowthistle (milk thistle)	Sonchus oleraceus
Corn Gromwell	Buglossoides arvensis
Cowvine	Ipomoea lonchophylla
Crassula	Crassula spp.
Deadnettle	Lamium amplexicaule
Dense-flower Fumitory	Fumaria densiflora
Dock	Rumex spp.
Doublegee (Spiny Emex)	Emex australis
Fat Hen	Chenopodium album
Fireweed	Senecio spp.
Fumitory	Fumaria spp.
Hedge Mustard	Sisymbrium officinale
Hexham Scent (King Island Melilot)	Melilotus indicus
Horehound	Marrubium vulgare
Hyssop Loosestrife	Lythrum hyssopifolia
Iceplant	Mesembryanthemum spp.
Indian Hedge Mustard	Sisymbrium orientale
London Rocket	Sisymbrium irio
Long Storks bill	Erodium botrys
Marshmallow	Malva parviflora

Common name	Scientific name
Mouse-eared Chickweed	Cerastium glomeratum
Night-scented Stock	Matthiola longipetala
Paterson's Curse	Echium plantagineum
Peppercress	Lepidium spp.
Prickly Lettuce	Lactuca serriola
Purple Goosefoot	Sclerobitum atriplicinum
Rough Poppy	Papaver hybridum
Saffron Thistle	Carthamus lanatus
Scarlet Pimpernel	Anagallis arvensis
Shepherd's Purse	Capsella bursa-pastoris
Skeleton Weed	Chondrilla juncea
Sorrel	Rumex spp.
Sternless Thistle	Onopordum acaulon
Toad Rush	Juncus bufonius Tree
Hogweed	Polygonum patulum
Turnip Weed	Rapistrum rugosum
Variegated Thistle	Silybum marianum
Vetch (Tares)	Vicia sativa
Volunteer Lupins	Lupinus spp.
Ward's Weed	Cardiaster annua
Wild Radish	Raphanus raphanistrum
Wild Turnip	Brassica tournefortii
Wreweed (Hogweed)	Polygonum aviculare

**GENERAL INSTRUCTIONS
RESISTANT WEEDS WARNING**

GROUP	FI	HERBICIDE
--------------	-----------	------------------

EZYCROP Diflufenican + MCPA Selective Herbicide is a member of the phenoxy and nicotinamide groups of herbicides and acts by inhibiting carotenoid biosynthesis at the phytoene desaturase step (PDS inhibitors) and disrupting plant cell growth. For weed resistance management EZYCROP Diflufenican + MCPA Selective Herbicide is both a Group F and a Group I herbicide. Some naturally occurring weed biotypes resistant to EZYCROP Diflufenican + MCPA Selective Herbicide and other Group F and I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by EZYCROP Diflufenican + MCPA Selective Herbicide or other Group F or Group I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, EZYCROP Pty Ltd accepts no liability for any losses that may result from the failure of EZYCROP Diflufenican + MCPA Selective Herbicide to control resistant weeds.

TOLERANCE

Some pre-emergence herbicides, such as atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Subterranean clover is particularly sensitive.

CEREALS

After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is not under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.

Warning (Oats): The tolerance of oat varieties Esk and Nile (the two main varieties grown in Tasmania) to EZYCROP Diflufenican + MCPA Selective Herbicide has not been tested. Test a small area of crop before using EZYCROP Diflufenican + MCPA Selective Herbicide over large areas. Consult your local EZYCROP representative for advice on specific varieties.

PASTURE

The tolerance of clover varieties to EZYCROP Diflufenican + MCPA Selective Herbicide can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress.

Warning: EZYCROP Diflufenican + MCPA Selective Herbicide may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter, particularly at rates in excess of 500 mL/ha and in areas of double spray. For this reason we recommend application prior to the 8 trifoliate leaf stage. However, at the lower rates (500 mL/ha and less) and under normal growing conditions, subsequent growth and seed yield should not be affected.

Under normal growing conditions, the following varieties have shown acceptable levels of foliage tolerance to EZYCROP Diflufenican + MCPA Selective Herbicide applied at 500 mL/ha:

ArrowLeaf: Zulu

Balansa: Paradana

Berseem: Sacromonte

Persian: Kyambro, Lupers, Maral

White: Haifa

Subterranean Clover: Daliak, Dalkeith, Denmark, Esperance, Geraldton, Goulburn, Karridale, Larissa, Leura, Mt.Barker, Nungarin, Rosedale, Seaton Park, Trikkala and Woogenellup.

The effects of this product on clover seed yield have been tested on the following varieties. Under normal growing conditions they show acceptable levels of tolerance to this product applied at 500 mL/ha:

Subterranean Clover: Esperance, Goulburn, Larissa, Seaton Park and Trikkala.

Warning: Rose and Strawberry clover have shown increased sensitivity to EZYCROP Diflufenican + MCPA Selective Herbicide. EZYCROP Diflufenican + MCPA Selective Herbicide may affect the seed yield of Subterranean Clover variety Woogenellup. Some pasture grasses, including Phalaris and Cocksfoot, may show some initial reduction in vegetative growth after application of EZYCROP Diflufenican + MCPA Selective Herbicide. Care should be exercised if sensitive clover varieties or grasses are included in the pasture sward. Varieties not listed should be tested before using EZYCROP Diflufenican + MCPA Selective Herbicide over large areas. Consult your local EZYCROP Pty Ltd representative for advice on specific varieties.

Subsequent Crops

To reduce effect on subsequent susceptible crops (e.g. canola), ensure thorough cultivation of soil prior to the sowing of these crops.

Mixing: To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly then add the remainder of the water. Agitate again before spraying commences. Reseal part-used product container immediately after use. Spray mixtures containing EZYCROP Diflufenican + MCPA Selective Herbicide should not be left to stand overnight. Prolonged periods of exposure to cold temperatures could result in settling out of the product in the mixture.

Warning: The rubber components present in some spraying units may be affected by exposure to the solvents in EZYCROP Diflufenican + MCPA Selective Herbicide and some other agricultural products. To reduce this risk it is recommended that the spray unit be thoroughly washed with a boom cleaner and fresh water after use. Contact the spray unit manufacturer to determine the suitability of the rubber components for use with agricultural products.

APPLICATION

Boom Sprayer: A minimum of 50 L of water per hectare should be used, however, for optimum results water rates of 70-100 L/ha are recommended. Increase the water volume if weed infestation is heavy or crop cover is dense. Complete coverage of weeds is essential.

Selective Herbicide

Aircraft (NSW, Vic, SA only): Apply in a minimum of 30 L water per hectare. Effective weed control will only be achieved where good coverage of leaf surface is achieved.

COMPATIBILITY

The following products are physically compatible with EZYCROP Diflufenican + MCPA Selective Herbicide as a two-way mixture in the spray tank but should only be used for the crops specified:

Crop	EZYCROP Diflufenican + MCPA Selective Herbicide	Compatible Product
Wheat, triticale and cereal rye only	Up to 750mL/ha	Hoegrass* (also barley), Tristar* Advance (1.5L rate only), Wildcat* (wild oats only)
Cereals (including undersown)	All rates	Chlorpyrifos (500g/L), dimethoate, Thiodan*.
Cereals (not undersown)	Up to 500mL/ha All rates	Ally*, Glean*, MCPA LVE, Logran* Bromoxynil (200g/L), 2,4-D Amine 500 Herbicide, Lontrel*, Tordon* 50-D, Cadence* (up to 115g only), Eclipse*
Wheat, barley, triticale, and cereal rye only (not undersown)		Achieve*
Wheat only (not undersown)		Topik*
Clover	Up to 750mL/ha	Targa*, Fusilade*
Subterranean Clover		Simazine (500g/L), simazine (500g/L) + paraquat (200g/L) mixture
	Up to 1 L/ha	2,4-DB amine (500 g/L)

When mixing with other herbicides, crop yellowing may be enhanced. When mixing with Hoegrass, Wildcat or Tristar Advance some reduction in the efficacy and speed of action of these products may occur. If the crop is

stressed, the application of the herbicide tank-mixtures may cause yield reduction. When mixing with Cadence a temporary wilting may be evident in some crops after application. Growers should seek advice before spraying recently released cereal varieties.

Use the recommended rates for both herbicides in the tank-mixture as well as the surfactant recommendation of the grass herbicide. If another herbicide is applied as a tank mix, observe the plantback restrictions on that label.

DO NOT add surfactant when mixing EZYCROP Diflufenican + MCPA Selective Herbicide and Ally.

Simazine: Refer to the simazine label for correct application rates, especially with regard to soil types. This product may be mixed in the spray tank with one of the following insecticides according to the directions for use on this product: Hallmark* 50EC, Dominex* 100EC, Karate*, Decis Options* and Talstar*.

Warning: DO NOT use crop oils with EZYCROP Diflufenican + MCPA Selective Herbicide or EZYCROP Diflufenican + MCPA Selective Herbicide tank mixtures with other products in cereals. As formulations of other manufacturer's products are beyond the control of EZYCROP Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF LIVESTOCK

Grazing Precaution: Sprayed weeds may become more palatable to stock and a higher intake of some weeds may result in stock poisoning and death from causes such as nitrate poisoning.

Care should be taken especially where Capeweed, Paterson's Curse and variegated thistles predominate in the pasture. Avoid grazing with young or breeding stock.

DO NOT graze horses or pigs on Paterson's Curse. If in doubt, contact your nearest Department of Agriculture.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish.

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area.

DO NOT store for prolonged periods in direct sunlight.

5L, 20L containers only:

Triple or preferably pressure rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Re-fillable containers:

If tamper evident seals are broken prior to initial use then the integrity of the contents cannot be assured.

Empty product as required into application equipment. Do not attempt to breach the valve system or filling point, or contaminate the container with water or other products. Ensure that the equipment used in transfer of the product is disconnected, triple rinsed and drained after each use.

When the container is empty, close all caps and valves and return the container to the point of purchase.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, DO NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the material safety data sheet (MSDS). A material safety data sheet for EZYCROP Diflufenican + MCPA Selective Herbicide is available from EZYCROP Pty Ltd on request.

CONDITIONS OF SALE

EZYCROP Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on EZYCROP Pty Ltd's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of EZYCROP Pty Ltd has any authority to add to or alter these conditions.

* Other trademarks.

BATCH NO:

DATE OF MANUFACTURE:

In an Emergency Dial 000 Police or Fire Brigade.