

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

EzyCrop Epoxiconazole 125 SC Fungicide

ACTIVE CONSTITUENT: 125 g/L EPOXICONAZOLE

GROUP	3	FUNGICIDE
--------------	----------	------------------

**For the control of leaf rust, stripe rust and septoria nodorum blotch
of wheat and leaf rust of barley, as specified in the
DIRECTIONS FOR USE table**

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

**Contents: 10L
(5L, 20L,)**

EzyCrop Pty Ltd
2/22 Horne Street
Elsternwick
VIC 3185
Australia

Tel: 03 9505 0044
ACN: 156 476 827



RLP
Approved



STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. Do not inhale vapours. If product on skin, immediately wash area with soap and water. Wash hands after use. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and half-piece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. After each day's use, wash gloves, respirator and if rubber, wash with warm water. Wash contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIALS SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS), which can be obtained from the supplier.

NOTICE TO BUYER

To the extent permitted by law, all conditions and warranties and statutory or other rights or action which buyer or any other user may have against EzyCrop Pty Ltd or Seller are hereby excluded. EzyCrop Pty Ltd hereby gives Notice to Buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by EzyCrop Pty Ltd or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. EzyCrop Pty Ltd liability shall, in all circumstances, be limited to replacement of the product or a refund of the purchase price paid therefor.

APVMA Approval No.

Batch No.

DOM

68670/58870

THIS PRODUCT IS NOT CONSIDERED TO BE A DANGEROUS GOOD UNDER THE AUSTRALIAN CODE FOR THE TRANSPORT OF DANGEROUS GOODS.

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

EzyCrop Epoxiconazole 125 SC Fungicide

ACTIVE CONSTITUENT: 125 g/L EPOXICONAZOLE

GROUP	3	FUNGICIDE
--------------	----------	------------------

**For the control of leaf rust, stripe rust and septoria nodorum blotch
of wheat and leaf rust of barley, as specified in the
DIRECTIONS FOR USE table**

IMPORTANT: READ THIS LEAFLET BEFORE USE

APVMA Approval No.

68670/58870

EzyCrop Pty Ltd
2/22 Horne Street
Elsternwick
VIC 3185
Australia

Tel: 03 9505 0044
ACN: 156 476 827

DIRECTIONS FOR USE

Crop	Disease	Rate per ha	Critical Comments
Wheat	Leaf rust (<i>Puccinia recondita</i>)	500 mL plus 200ml non-ionic surfactant per 100L spray-mix	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59). DO NOT apply later than ZGS 59. Repeat spraying may be required for stripe rust. Regularly monitor the crop from 3-4 weeks after spraying for signs of re-infection and treat accordingly.
	Stripe rust (<i>Puccinia striiformis</i>)		
	Septoria nodorum blotch (<i>Septoria nodorum</i>)		
Barley	Leaf rust (<i>Puccinia hordei</i>)	250 to 500 mL plus 200ml non-ionic surfactant per 100L spray-mix	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and ear emergence complete (ZGS 32-59). DO NOT apply later than ZGS 59. Apply the higher rate when disease is present on the top leaf or conditions are favourable for disease development at these stages of crop development.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

WITHHOLDING PERIODS

DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION.

DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 6 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

EzyCrop Epoxiconazole 125 SC is a broad-spectrum contact and systemic fungicide. It is taken up via the stem and foliage and is translocated upwards and outwards to provide protection to new growth. Thorough coverage of the crop is necessary for best results.

Check the crop regularly for any symptoms of disease infection. Apply as soon as disease is present on the flag leaf or flag-2. Application beyond flag leaf emergence (ZGS39) may not provide any yield benefit, particularly under dry seasonal conditions.

MIXING

EzyCrop Epoxiconazole 125 SC is a suspension concentrate formulation. Add the product to the half-filled spray tank while agitating. Continue to agitate while topping up the tank and during spraying.

APPLICATIONGround Application

Apply in a water volume of between 50 and 100L/ha, using flat fan nozzles operating at around 50 cm above the top of the crop. Use the higher water volume in crops with heavier canopies.

Aerial Application

Apply with suitable aircraft set up and operated to apply fungicides to cereal crops in a minimum water volume of 20L/ha.

COMPATIBILITY

For information on compatibility please contact your local reseller or EzyCrop Pty Ltd representative.

FUNGICIDE RESISTANCE WARNING

GROUP	3	FUNGICIDE
-------	----------	-----------

EzyCrop Epoxiconazole 125 SC is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides

are used repeatedly. These resistant fungi will not be controlled by this product or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, EzyCrop Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

RE-ENTRY PERIOD

DO NOT allow entry into treated areas until the spray has dried.

CAUTION Export of treated cereals

Growers should note that Maximum Residue Limits (MRLs) or import tolerances do not exist in all markets for cereals treated with EzyCrop Epoxiconazole 125 SC Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing cereals for export, please check with EzyCrop Pty Ltd or your grain exporter for the latest information on MRLs and import tolerances BEFORE using EzyCrop Epoxiconazole 125 SC Fungicide.

LIVESTOCK DESTINED FOR EXPORT MARKETS

The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.

EXPORT SLAUGHTER INTERVAL (ESI):

AFTER OBSERVING THE PERIOD FOR GRAZING OR CUTTING FOR STOCKFOOD, LIVESTOCK THAT HAVE BEEN GRAZED ON OR FED TREATED CROPS SHOULD BE PLACED ON CLEAN FEED FOR 3 DAYS PRIOR TO SLAUGHTER.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Highly toxic to aquatic organisms. Do NOT apply under weather conditions or from spraying equipment which could be expected to cause spray drift onto adjacent crops, croplands, livestock, pastures, wetlands, waterbodies or watercourses. Do NOT apply EzyCrop Epoxiconazole 125 SC Fungicide aerially if waterbodies, watercourses or wetlands are within 30 metres downwind of the application area. Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area out of direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should NOT be burnt.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Will irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. Do not inhale vapours. If product on skin, immediately wash area with soap and water. Wash hands after use. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and half-piece respirator. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. After each day's use, wash gloves, respirator and if rubber, wash with warm water. Wash contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIALS SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet (MSDS), which can be obtained from the supplier.

NOTICE TO BUYER

To the extent permitted by law, all conditions and warranties and statutory or other rights or action which buyer or any other user may have against EzyCrop Pty Ltd or Seller are hereby excluded. EzyCrop Pty Ltd hereby gives Notice to Buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by EzyCrop Pty Ltd or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. EzyCrop Pty Ltd liability shall, in all circumstances, be limited to replacement of the product or a refund of the purchase price paid therefor.