

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

EzyCrop Imidacloprid 200 SC Insecticide

ACTIVE CONSTITUENT: 200g/L IMIDACLOPRID

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

**For the control of various insect pests of certain vegetables and turf
as specified in the Directions for Use table**

**IMPORTANT: READ THIS BOOKLET BEFORE OPENING OR USING
THE PRODUCT**

Contents 1L
(5L, 10L, 20L)

EzyCrop Pty Ltd
2/22 Horne Street
Elsternwick
VIC 3185
Australia

Tel: 03 9505 0044
ACN: 156 476 827



STORAGE & DISPOSAL

Store in the closed, original containers in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, and or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS), which is available from the supplier or from EzyCrop Pty Ltd.

NOTICE TO BUYER

EzyCrop Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence, use under abnormal conditions or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on EzyCrop Pty Ltd's skill or judgment in purchasing or using the product and every person dealing with this product does so at their own risk

APVMA Approval No; 68673/58873

Batch No:

Date of Manufacture:



CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

EzyCrop Imidacloprid 200 SC Insecticide

ACTIVE CONSTITUENT: 200g/L IMIDACLOPRID

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

**For the control of various insect pests of certain vegetables and turf
as specified in the Directions for Use table**

**IMPORTANT: READ THIS BOOKLET BEFORE OPENING OR USING
THE PRODUCT**

APVMA Approval No: 68673/58873

EzyCrop Pty Ltd
2/22 Horne Street
Elsternwick
VIC 3185
Australia

Tel: 03 9505 0044
ACN: 156 476 827

DIRECTIONS FOR USE**FOLIAR SPRAY APPLICATIONS**

Crop	Insect	Rate	WHP	Critical Comments
Cucurbits	Green Peach Aphid	25mL/100L or 300mL/ha	1 days	Apply at first sign of aphid infestation.
Capsicum			7 days	
Eggplant				
Potato				
Tomato	Grey cabbage Aphid Turnip Aphid		3 days	Apply at first sign of aphid infestation. Add a wetting agent.
Brassicas			7 days	
Sweet potato	Silverleaf Whitefly including type B	25mL/100L or 250mL/ha	7 days	Apply at first sign of whitefly or melon thrips infestation. Apply dilute sprays (25mL/100L) to run off. Ensure thorough coverage of underside of leaves. Use of droppers will improve coverage of underside of leaves.
Cucumber			1 day	
Egg-plant	Melon Thrips		7 days	
Turf	First instar larvae of: African Black Beetle, Argentinian Scarab, Pruinose Scarab	2.5L/ha or 25mL/100m ² Spray with at least 400 L water per hectare to ensure even coverage. Preferably spray on to wet or dewy grass. Irrigate with 12 mm of water commencing within one hour of application.	-	Apply at peak egg hatch that is mid Spring to mid Summer depending on species.
	Larvae of Billbug			Monitor adult activity through late Spring and early Summer. Spray when numbers peak, or when small larvae (4mm) are found in the thatch or surface soil. Early application is essential to minimise grass damage due to feeding.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

Cucurbits: DO NOT harvest for 1 day after application

Tomatoes: DO NOT harvest for 3 days after application

Brassicas, Capsicum, Egg Plant, Potatoes, Sweet potatoes: DO NOT harvest for 7 days after application

DO NOT graze any treated area, including turf, or cut for stock food.

DO NOT feed produce or turf clippings from treated area to animals, including poultry.

GENERAL INSTRUCTIONS**INSECTIDE RESISTANCE WARNING**

GROUP	4A	INSECTICIDE
-------	----	-------------

For insect resistance management EzyCrop Imidacloprid 200 SC Insecticide is a Group 4A Insecticide.

Some naturally occurring insect biotypes resistant to EzyCrop Imidacloprid 200 SC Insecticide and other group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if EzyCrop Imidacloprid 200 SC

Insecticide or other Group 4A insecticides are used repeatedly. The effectiveness of EzyCrop Imidacloprid 200 SC Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, EzyCrop Pty Ltd accepts no liability for any losses that may result from the failure of EzyCrop Imidacloprid 200 SC Insecticide to control resistant insects. EzyCrop Imidacloprid 200 SC Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, EzyCrop Pty Ltd representative or local agricultural department agronomist.

RESISTANCE MANAGEMENT STRATEGY

Aphids, whitefly and melon thrips in various crops

Do not apply EzyCrop Imidacloprid 200 SC Insecticide (or other Group 4A insecticides) in consecutive sprays within and between seasons. Rotate with registered insecticides from other mode of action groups.

MIXING/APPLICATION

Prior to pouring, shake container vigorously, then add the required amount of EzyCrop Imidacloprid 200 SC Insecticide to water in the spray vat while stirring or with agitators in motion.

EXPORT OF TREATED PRODUCE

Growers' should note that MRLs or import tolerances do not exist in all markets for edible produce treated with EzyCrop Imidacloprid 200 SC Insecticide. If you are growing edible produce for export, please check with EzyCrop Pty Ltd for the latest information on MRLs and import tolerances before using EzyCrop Imidacloprid 200 SC Insecticide.

COMPATIBILITY

EzyCrop Imidacloprid 200 SC Insecticide is compatible with propineb, bitertanol or methamidophos. Do not mix concentrates together but add each to the spray tank separately. As formulations of other manufactures' products are beyond the control of EzyCrop Pty Ltd, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, EzyCrop Pty Ltd cannot be responsible for the behaviour of such mixtures.

PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. DO NOT graze any treated area, or cut for stock food. DO NOT feed produce or turf clippings from treated area to animals, including poultry.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways and drains with the chemical or used containers.

STORAGE & DISPOSAL

Store in the closed, original containers in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, and or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When preparing product for use wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. Wash hands after use. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

For further information, refer to the Material Safety Data Sheet (MSDS), which is available from the supplier or from EzyCrop Pty Ltd.

NOTICE TO BUYER

EzyCrop Pty Ltd shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence, use under abnormal conditions or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on EzyCrop Pty Ltd's skill or judgment in purchasing or using the product and every person dealing with this product does so at their own risk