

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

EzyCrop MCPA 500 Selective Herbicide

ACTIVE CONSTITUENT: 500 g/L MCPA present as the dimethylamine salt

GROUP **I** HERBICIDE

For the selective control of broadleaf weeds in cereals, linseed, pastures, sugar cane and turf as per Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.

Contents: 5 L (20L, 200L, 1000L)

EzyCrop Pty Ltd
2/22 Horne Street
Elsternwick
VIC 3185
Australia

Tel: 03 9505 0044
ACN: 156 476 827



RLP
Approved



GENERAL INSTRUCTIONS

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours following application.

STORAGE AND DISPOSAL**(5L, 20L, 200L only):**

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and products should not be burnt.

Refillable containers (1000L only): Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin when opening the container and preparing spray and using the prepared spray. Wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product on skin immediately wash area with soap and water. If product or spray in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE

"EzyCrop Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on EzyCrop Pty Ltd's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of EzyCrop Pty Ltd has any authority to add to or alter these conditions."

APVMA Approval No.: 68816 / 59149

Batch No.:

Date of Manufacture:

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

EzyCrop MCPA 500 Selective Herbicide

ACTIVE CONSTITUENT: 500 g/L MCPA present as the dimethylamine salt

| | | |
|-------|---|-----------|
| GROUP | I | HERBICIDE |
|-------|---|-----------|

For the selective control of broadleaf weeds in cereals, linseed, pastures, sugar cane and turf as per Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT.

Contents: 5 L (20L, 200L, 1000L)

APVMA Approval No: 68816 / 59149

EzyCrop Pty Ltd
2/22 Horne Street
Elsternwick
VIC 3185
Australia

Tel: 03 9505 0044
ACN: 156 476 827

DIRECTIONS FOR USE:

RESTRAINTS: DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.
DO NOT apply if rain is likely within 6 hours.

1. Field Crops

| Situation and Crop | Weeds | State | Rate | Critical Comments |
|--------------------------------|---|-----------------------------|--|---|
| Wheat, Barley, Oats, Triticale | Refer to Weed Table | SA, Tas only | 700ml – 2.1L/ha | Apply through the tillering stage after the crop plants have 5 leaves and before they commence jointing. |
| | | Qld only | | Apply from early tillering (4-5 fully emerged main stem leaves plus one or more tillers) until before the boot stage (visible swelling of the head at the top of the main stem). |
| | | NSW, ACT only | | Apply 700ml rate after crop has reached 5 leaf stage and weeds marked E in weed table have 3-6 leaves. Higher rates after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller. |
| | | Vic only | 700ml – 1.5L/ha | Apply 700ml rate after crop has reached 2-3 leaf stage and weeds marked E in weed table have 2-4 leaves. Higher rate after the crop plants have 5 leaves to fully tillered. |
| | | WA only | 900ml - 2L/ha | Apply to crop from 5 leaf to Flag leaf emergence. (Zadoks 15-37) |
| Wheat, Barley, Oats, Triticale | Soldier Thistle | SA only | 700ml plus 200ml/ha EzyCrop Clopyralid 750 | Apply to crop at tillering stage after the crop has 5 leaves and before the crop commences jointing. |
| | Skeleton Weed | | 1L plus 500ml/ha EzyCrop Clopyralid 750 | Apply to rosettes from 5-15cm diameter when the crop is between the 5 leaf and late tillering growth stages. It is essential that all the Skeleton weed is emerged and minimum of 5cm diameter which is usually not before August. This rate will provide control of Skeleton Weed until harvest and will substantially reduce seed numbers the following season. |
| Linseed | Refer to Weed Table | Qld only | 700ml – 2.1L/ha | Apply when crop is 10-15cm tall with at least 170L/ha of water. Do not spray after buds appear. Some wilting effect may be seen following application but crop usually recovers quickly. Do not apply by aircraft. |
| | | NSW, ACT, Vic, SA, Tas only | 700ml – 1.1L/ha | |
| Sugar Cane | Blue Top, Chinese Burr, Flannel Weed, Gambia Pea, Bell Vine, Streaked Rattle Pod, Bindweed, Pink Convolvulus, Cupids Flower, Merremia Vine, Morning Glory | QLD only | 1.4L/ha | POST-EMERGENT: Apply as directed inter-row spray. Do not exceed rates with phenoxy sensitive varieties. |
| | As above plus Fat Hen, Noogoora Burr | | 2.1L/ha | |
| | All of the above | | 10L/ha | |
| Flax | Refer to Weed Table | Vic, Tas only | 490ml/ha | Apply to crop when 8-20cm high |

1. FIELD CROPS(continued)

| Situation and Crops | Weeds | State | Weed Stage | Rate | Critical Comments |
|---------------------|--|------------------------------|------------|--|--|
| Field Peas | Prickly Lettuce (<i>Lactuca serriola</i>) | NSW, Vic, Tas, SA only | 4 leaf | 125ml plus 125mls EzyCrop Diflufenican500 | <p>FIELD PEAS: Apply early post-emergence after the third node stage and before the start of flowering.</p> <p>Warning: Field peas grown on high pH soils in the presence of free lime may be less tolerant to Agility.</p> <p>APPLICATION AND WEED CONTROL Apply when weeds are actively growing. For optimum results apply 4 to 6 weeks post-sowing.</p> <p>Application beyond 8 weeks post-sowing may result in reduced levels of weed control. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as:</p> <ul style="list-style-type: none"> • high crop and weed density, • late season germinations, • abnormal weed growth (including early flowering); higher rates of product (up to the maximum rate of application specified for that weed) may be required. <p>EzyCrop Diflufenican 500 will not effectively control:</p> <ul style="list-style-type: none"> • Regrowth of suppressed weeds • Transplanted weeds • Regrowth from rhizomes or roots • Weeds growing under stress from previous herbicide applications |
| | | | 6 leaf | 150ml plus 150mls EzyCrop Diflufenican500 | |
| | Charlock (wild Mustard) (<i>Sinapis arvensis</i>) | | 4 leaf | 125ml plus 125mls EzyCrop Diflufenican500 | |
| | | | 6 leaf | 150ml plus 150mls EzyCrop Diflufenican500 | |

2. PASTURES, SEED CROPS & TURF

| Situation & Crop | Weeds | State | Rate | Critical Comments |
|----------------------------------|--|----------------------------|-------------------------------------|---|
| Grass Pastures | Refer to Weed Table | All states | 700ml – 4L/ha | Apply to established grass-based pastures. Damage may occur to Legumes if present. See "Legume Tolerance" below. NOTE: Capeweed, paterson's curse and variegated thistle may be poisonous to stock after spraying. |
| | Soldier Thistle | SA only | 700ml – 2.1L/ha | Use low rate on seedlings with no more than 2 true leaves. Use 100 litres of water for best results. If clover is present, do not exceed 600ml. |
| | Seedling and established Horehound | NSW, ACT only | 3L/ha | Apply in autumn when Horehound plants are at seedling stage. If sub-clover or medic plants are present they will be damaged by this application. |
| Grass Seed Crops | Refer to Weed Table | Qld, SA, Tas, Vic, WA only | 700ml – 4L/ha | Apply to established grass-seed crops |
| Sub-clover seed crops | Refer to Weed Table | Vic, SA only | 700mL-1.5L/ha | Do not exceed rate specified. Spray in Autumn and early Winter. |
| Pastures - Spray/Graze Technique | Caltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherds Purse, Saffron, Slender, Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip | NSW, ACT, Vic only | 350mL-1.4L/ha | Spray actively growing 6-8 weeks old weeds. Introduce stock 7-10 days after spraying, preferably sheep as cattle are less effective. Use stocking rate at least 5 times heavier than normal until weeds have been reduced but before survival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed short before spraying. Refer "Legume Tolerance". PRECAUTION: An increased quantity of poisonous plants may be eaten by stock using Spray/Graze, eg. caltrop, capeweed, paterson's curse and variegated thistle and deaths could result from causes such as nitrate poisoning. With paterson's curse preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. Do not graze horses or pigs on paterson's curse. Observe Withholding Period of at least 7 days after spraying. |
| | Refer to Weed Table | Tas only | 700mL/ha | |
| | Capeweed, Dandelion, Seedling Dock, Three Corned Jack, Geranium, Erodium, Mustard, Annual Thistles, Turnip Weed, Wild Radish, Wild Turnip | WA only | 925ml-1.85L/ha | |
| | Soldier Thistle | SA only | 500mL/ha | |
| Turf | Refer to Weed Table | All States | 500mL/ha or 5mL per 100 sq.m | Newly laid - Delay until well established or at least two months old. Apply in high volume to actively growing weeds. See "Legume Tolerance". Buffalo grass is more easily damaged than other grasses. |
| | | | 1.4-2.8L/ha or 14-28mL per 100 sq.m | Established - Apply in high volume to actively growing weeds. Do not mow for 2 days before application. Some transitory damage may occur to fine turf grasses and strawberry clover. See "Legume Tolerance". Buffalo grass is more easily damaged than other grasses. |

3. SPOT SPRAYING

| Situation & Crop | Weeds | State | Mixing Rates/Critical Comments |
|----------------------|---------------------|------------|--|
| High Volume Spraying | Refer to Weed Table | All States | Add 1/10th of rate on Weed Table to 150 litres of water. Each 150 litres of mix will cover 1000m ² (1/10th ha.) eg. If rate in Weed Table is 1.4L use 140mL/150L water. |
| Knapsack Application | | | Add 1/100th of rate on Weed Table to 10 litres of water. Each 10 litres of mix will cover 100m ² (1/100th ha.) eg. If rate in Weed Table is 1.4L use 14mL/10L water. |

4. Non Crop Areas, Fallow Land, Industrial & Commercial Areas, Rights-of-Way

| Situation & Crop | Weeds | State | Rate | Critical Comments |
|---|---------------------|------------|---------------------|--|
| Non Crop Areas, Fallow Land, Industrial & Commercial Areas, Rights of Way | Refer to Weed Table | All States | Refer to Weed Table | Apply at rates shown in Weed Table when weeds only are present and no crop or pasture is involved. |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: CEREALS, PASTURE – DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 7 DAYS AFTER APPLICATION.

HARVEST WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours of application.

APPLICATION/MIXING INFORMATION

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

BOOM SPRAYING - Use 30-120 litres water per hectare. - Use minimum of 170 litres water per hectare for LINSEED.

AERIAL SPRAYING - Use 10-90 litres water per hectare.

RESISTANT WEEDS WARNING

| | | |
|--------------|----------|------------------|
| GROUP | I | HERBICIDE |
|--------------|----------|------------------|

EzyCrop MCPA 500 Selective Herbicide is a member of the Phenoxy group of herbicides. EzyCrop MCPA 500 has the Disruptors of plant cell growth mode of action. For weed resistance management EzyCrop MCPA 500 is a Group I herbicide.

Some naturally-occurring weed biotypes resistant to EzyCrop MCPA 500 and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by EzyCrop MCPA 500 or other disruptors of plant cell growth herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, EzyCrop Pty Ltd accepts no liability for any losses that may result from the failure of EzyCrop MCPA 500 to control resistant weeds.

EQUIPMENT MAINTENANCE AND USAGE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been cleaned with Spraymate Tank & Equipment Cleaner or well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses.

COMPATIBILITY/TANK MIXES

This product can be tank mixed with EzyCrop Diflufenican, Bromoxynil, Clopyralid, 2,4-D/Picloram, and metsulfuron-methyl products. It can also be tank mixed with Kamba®, Sertin®, Flowable Terbutryn®, Tribunil® Imidan®, and Sprayseed®. The effectiveness of Imidan®, Le Mat®/Comrad, Metasystox® and Dimethoate may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations the addition of a soya phospholipid-propionic acid adjuvant is recommended. When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds. DO NOT spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result.

Drift Warning: DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, fruit trees, vegetables, legume crops and pastures, oilseed crops and susceptible trees (eg. kurrajongs, belahs, eucalypts).

Legume Tolerance: The use of MCPA is preferred to 2,4-D Amine or Ester in situations where legumes are undersown in the crop.

NB: The tolerance of pasture legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. DO NOT spray cereal crops undersown with lucerne as damage may result.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the product or used container.

STORAGE AND DISPOSAL

(5L, 20L, 200L only):

Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and products should not be burnt.

Refillable containers (1000L only): Store in the closed, original container in a dry, cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate the skin. Avoid contact with eyes and skin when opening the container and preparing spray and using the prepared spray. Wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product on skin immediately wash area with soap and water. If product or spray in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone 13 11 26).

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS).

CONDITIONS OF SALE

EzyCrop Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on EzyCrop Pty Ltd's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of EzyCrop Pty Ltd has any authority to add to or alter these conditions."

® - Other commercial products.

WEED TABLE:

Where weeds are to be sprayed in a crop or pasture use only the rates given for particular crop or situation indicated under the Directions for Use table.

In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage. This product should only be used in those states where a rate or range of rates is indicated for the particular weeds listed.

APPLICATION RATE PER HECTARE

*E: refer to Critical Comments for Wheat, Barley, Oats, Triticale

| WEEDS CONTROLLED | Vic | NSW, ACT | SA | Tas | Qld | WA | Critical Comments |
|---------------------------|----------|------------|-------|--------|--------|-------|--|
| Ball Mustard | - | - | 1.7L | - | - | - | Apply at rosette stage. |
| Bathurst Burr | 2L | 1-2L | 2L | 2L | 2L | 2L | Spray young seedlings only. |
| Black Bindweed | - | 1.4-2L | - | - | 1.4-2L | - | Apply at young stage prior to flowering |
| Californian Burr | - | 1-2L | - | 2.2L | - | - | Spray young seedlings only. |
| Cape Tulip | - | - | - | - | - | 2L | Apply just prior to flowering. Gives mild suppression only. |
| Capeweed | - | 2.1L | 2.1L | 2.1L | 2.1L | - | Apply at seedling stage. |
| Carrot Weed | 2L | - | - | - | 2L | - | Apply at seedling stage. |
| Charlock | 700mL *E | 1L *E | 700mL | 1-2.2L | 700mL | - | Spray up to rosette stage. |
| Cutleaf Mignonette | - | - | 2.1L | - | - | - | Apply at young rosette stage. |
| Dandelion | 2L | 2L | 2L | 2L | 2L | 2L | Apply at young rosette stage. |
| Deadnettle | - | 2.1L | - | - | 2.1L | - | Moderately susceptible. Apply at seedling stage. |
| Docks | 1.5L | - | 2.1L | 2.1L | - | - | Apply at rosette stage and before flower heads emerge. Only the top growth of old established plants will be affected. |
| Fat Hen | 2L | 1-2L | - | 2L | 2L | - | Spray up to rosette stage. |
| Fennel | 2.7L | 2.7L | 2.7L | 2.7L | 2.7L | - | Apply at young stage. |
| Field Bindweed | 2L | - | 2L | 2L | 2L | 2L | Apply at bud stage, repeating over several seasons. |
| Field Cress | - | - | - | 2.2L | - | - | Apply at young stage. |
| Flatweed or Catsear | - | - | - | 2.2L | - | 1.4L | Apply when weeds are 10-15cm high. |
| Fumitory (red) | 1.4L | 1.4L | 1.4L | - | 1.4L | - | Apply at young stage prior to flowering. |
| Hoary Cress | 4.5L | - | 2.1L | - | 4.5L | - | Apply before flowering. |
| Horehound | - | 3L | 2.1L | - | - | - | Apply at rosette stage. |
| Hedge Mustard | 1.1L *E | - | 700mL | 700mL | 700mL | 1.25L | Apply at rosette stage. |
| Hexham Scent or Melilotus | 1.5L | 1.2L | 1.4L | - | 1.4L | - | Apply at young seedling stage. |
| Lesser Swinecress | - | - | - | 2.2L | - | - | Apply at young stage. |
| Lincoln Weed | - | - | 700mL | - | - | - | Apply at rosette stage. |
| London Rocket | - | - | - | - | - | 1.25L | |
| Lupins | - | 700mL-1.5L | - | - | - | 1.25L | Spray up to 10cm high. |
| Mintweed | 2L | 2L | - | - | 2L | - | Spray young seedlings only. |

APPLICATION RATE PER HECTARE continued

| WEEDS CONTROLLED | Vic | NSW, ACT | SA | Tas | Qld | WA | Critical Comments |
|-----------------------|-----------|------------|------------|----------|------------|----------|--|
| Mustards | 1.1L *E | 1L *E | 1.1L | 1.1L | - | 1.25L | Spray up to rosette stage. |
| Noogoora Burr | 2L | 1-2L | 2L | - | 2L | 2L | Spray young seedlings only. |
| Opium Poppy | - | - | - | 2.2L | - | - | Apply at young stage. |
| Paterson's Curse | - | 1-1.5L | - | - | 4L | - | Apply early rosette stage. |
| Pimpernel | - | - | 700mL | - | - | - | Apply at rosette stage. |
| Plantains | - | 2L | - | 2L | 2L | - | Apply at young stage. |
| Rapeseed | - | 1-1.5L | - | - | - | - | Spray up to rosette stage. |
| Rough Poppy | - | 700mL-1.5L | - | - | - | - | Spray up to rosette stage. |
| Safflower | - | 700mL-1L | - | - | - | - | Spray up to rosette stage. |
| Shepherd's Purse | - | - | - | 2.2L | - | - | Apply at rosette stage. |
| Skeleton Weed | - | 1.5-2L | - | - | - | - | Spray rosettes before aerial growth commences. |
| Sorrel | - | - | 2.1L | - | - | - | Apply to rosette stage. |
| Stinkwort | 2L | 1-1.5L | 2L | - | - | 2L | Apply when weeds are 10-15cm high. |
| Sunflower (seedlings) | - | 1-1.5L | - | - | - | - | Spray up to rosette stage. |
| Thistles: | | | | | | | |
| - Artichoke | 1.6L | - | 2.1L | - | 2L | 700mL | Apply at rosette stage. |
| - Nodding | - | 2.5L | - | - | - | - | Spray early rosettes. Repeat spray required in following Spring or Autumn. |
| - Saffron | 1.5L-2.5L | 1-2L | 1.4-2.5L | 1.4-2.5L | 1.4-2.5L | 1.5-2.5L | Spray up to rosette stage. (September WA). |
| - Scotch or Cotton | 2.2L | - | 700mL-1.4L | 2.2L | - | - | Apply at seedling stage. |
| - Slender | 3L | 1-2L | 1L | 3L | - | 3L | Spray young seedlings only. |
| - Soldier | 1.1L | - | 1.1L | - | - | - | Spray young seedlings. Use higher rate for larger weeds. |
| - Spear | - | 1.5-2L | - | 2.2L | - | - | Apply at rosette stage. |
| - Star | - | - | 1-1.4L | - | - | - | Apply at seedling stage. Use higher rate for larger weeds. |
| - Variegated | 1.5L | 1-2L | 700mL-1.4L | 1.5L | 1.5L | 1.5L | Spray young seedlings only. Use 2L/ha (SA) for well developed weeds. |
| Turnip Weed | 1.1L *E | 1L *E | 1L | - | - | - | Spray up to rosette stage. |
| Wild Artichoke | - | - | 2.1L | - | - | - | Apply at rosette stage. |
| Wild Radish | 1.4L | 1L *E | 1.4L | 2L | 700mL-1.1L | 1.2L | Spray up to rosette stage. |
| Wild Turnip | 1.1L *E | 1L *E | 700mL | 2L | 700mL-1.1L | 900mL | Spray up to rosette stage. |