

Company Name: EZYCROP PTY LTD
Product Name: Ezycrop Phosphorous Systemic Fungicide
APVMA Approval No: 81494/103211



Label Name:	Ezycrop Phosphorous Systemic Fungicide
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	ACTIVE CONSTITUENT: 600 g/L PHOSPHOROUS (PHOSPHONIC) ACID PRESENT AS MONO (AND) DI POTASSIUM PHOSPHITE
Mode of Action:	GROUP 33 FUNGICIDE
Statement of Claims:	Effective against Phytophthora diseases and Downy Mildew in various situations as per Directions for Use Table.
Net Contents:	1 L 5 L 20 L 200 L 1000 L
Restraints:	RESTRAINT: DO NOT apply this product at volumes which cause excessive run off.
Directions for Use:	This section contains file attachment. File Name: Ezy Phos- DFU - 0715.doc File Size: 51712 bytes
Other Limitations:	

Withholding Periods:	<p>WITHHOLDING PERIOD: Subterranean Clover: DO NOT GRAZE OR CUT FOR STOCKFEED FOR 14 DAYS AFTER APPLICATION.</p> <p>Citrus, Cucurbits, Avocado, Grapes, Pineapples: NOT REQUIRED WHEN USED AS DIRECTED.</p>
-----------------------------	---

Trade Advice:	
----------------------	--

General Instructions:	<p>This section contains file attachment. File Name: Ezy Phos- GI - 0915.docx File Size: 19937 bytes</p>
------------------------------	--

Resistance Warning:	<p>FUNGICIDE RESISTANCE WARNING GROUP 33 FUNGICIDE</p> <p>For fungicide resistance management, Ezycrop Phosphorous Systemic Fungicide is a Group 33 fungicide. Some naturally occurring individual fungi resistant to Ezycrop Phosphorous Systemic Fungicide and other Group 33 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Ezycrop Phosphorous Systemic Fungicide, and other Group 33 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Ezycrop Pty Ltd accepts no liability for any losses that may result from the failure of Ezycrop Phosphorous Systemic Fungicide to control resistant fungi.</p>
----------------------------	--

Precautions:	
---------------------	--

Protections:	<p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT This product is harmful to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.</p>
---------------------	---

Storage and Disposal:	<p>STORAGE AND DISPOSAL Store in the closed, original container in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Spillage should not be directed to drains but absorbed in sawdust and the absorbent material disposed of in a sealed container at an approved disposal site. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.</p> <p>For refillable containers: Empty contents fully into application equipment. Close all valves and return to designated collection point for refill or storage.</p>
------------------------------	---

Safety Directions:	<p>SAFETY DIRECTIONS</p> <p>May irritate eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves, goggles and disposable face mask. If clothing becomes contaminated with product or wet with spray, remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing. Wash hands after use.</p>
---------------------------	--

First Aid Instructions:	<p>FIRST AID</p> <p>If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.</p>
--------------------------------	---

First Aid Warnings:	
----------------------------	--

DIRECTIONS FOR USE

TREE AND VINE CROPS

CROP	DISEASE	STATE	TREATMENT METHOD	RATE	CRITICAL COMMENTS
AVOCADO	Phytophthora Root Rot (<i>Phytophthora cinnamomi</i>)	QLD, NSW, SA, VIC, WA, ACT	Injection	Trunk Injection, Skeletal trees: 1 st year 5 mL undiluted product per metre of canopy diameter. Other situations: 2.5 mL product diluted with 7.5 mL water per metre of canopy diameter.	Inject trees at spring flush maturity - Repeat treatment in February or March. Drill holes 5 mm in diameter and 25 – 50 mm deep with slight downward angle in trunk. Syringes should be placed in the main trunk of the tree and spaced evenly around the circumference of the trunk. Suitable for use with Chemjet tree injectors, Ag-murf gun or hydraulic tree injection. Do not prune trees back before injection process as burning of new growth may occur. Do not inject trees in winter months. Do not cut back the canopy of injected trees. Do not add anything other than water to Ezycrop Phosphorous Systemic Fungicide for trunk injection. DO not inject more liquid in a lesser number of syringes than directed. Solution concentrate: 165 - 200 mL/100 L. Amount per unit: Spray to run off – 7.5 to 10 L per adult tree. 1.Curative: Apply every 3 weeks until disease is under control. 2.Preventative: Apply every 5 - 6 weeks.
YOUNG OR SMALL CITRUS.	Phytophthora Root Rot (<i>P. nicotianae</i>), and Collar Rot (<i>P. citrophthora</i>)	All states	Foliar spray	170 - 330 mL/ 100 L to leaf wetness (by boom or similar high volume sprayer)	Two applications <i>1st Application:</i> Late winter prior to flowering. <i>2nd Application:</i> Autumn applied to mature fruit. Repeat applications annually to maintain protection for the plant. Depending on condition at time of application use a non-ionic wetting agent in conjunction with Ezycrop Phosphorous Systemic Fungicide. WHERE DISEASE INCIDENCE IS HIGH OR WELL ESTABLISHED. For effective control, apply as a protectant before above ground symptoms of decline and collar rot become evident, spray trees for even coverage. Do not apply under high temperatures (above 35°C) particularly if humidity is low, or to moisture stressed trees. LOW PHYTOPHTHORA PRESSURE, WELL DRAINED SOIL Removal of fruit from affected plants will enhance recovery. WARNING: Soft skin citrus trees may develop leaf burn and growth retardation following foliar application or soil drench of Potassium Phosphonate at rates recommended for established trees. <div style="border: 1px solid black; padding: 5px; margin-top: 10px;"> WARNING: 1. Foliar application to fruit may cause damage to fruit. 2. Soft skinned citrus trees may develop leaf burn and growth retardation following foliar application or soil drench. Before applying this product for the first time, a small test area should be sprayed and observed for at least 1 week. </div>
MATURE CITRUS		All states	Foliar spray	13.3 L/ha in 3000 - 4000 L of water. 53 mL/12 L per tree. 8.3 L/ha in 2000 - 5000 L of water. 33 mL/12 L per tree	

CROP	DISEASE	STATE	TREATMENT METHOD	RATE	CRITICAL COMMENTS
GRAPES	Downy Mildew (<i>Plasmopara viticola</i>)	All States	Foliar spray	2 L/ha early season, small canopies 2.7 - 4 L/ha Mid-late season, large canopies	It is essential that the rate of Ezycrop Phosphorous Systemic Fungicide is adjusted to the vine-row volume (i.e. the volume of vine foliage per hectare). An application volume of 500 L/ha is suggested at the start of the season, increasing to 2000 L/ha in a vigorous crop at full canopy. Spray timing is critical. For best results apply Ezycrop Phosphorous Systemic Fungicide as a tank mix with protectant fungicides such as Mancozeb/Dithane, Copper Oxychloride etc, to ensure both pre- and post-infection activity. Ezycrop Phosphorous Systemic Fungicide should be applied at times of high disease risk, especially between the time that conditions are conducive to Downy Mildew infection and the appearance of oil spots. Ensure spray coverage is adequate and that the appropriate rate of Ezycrop Phosphorous Systemic Fungicide is applied to match vine growth, particularly from mid-season onwards and especially where grapes are grown on root stock.

NON TREE AND VINE CROPS

CROP	DISEASE	STATE	TREATMENT METHOD	RATE	CRITICAL COMMENTS
CUCURBITS	Downy Mildew (<i>Pseudoperonospora cubensis</i>)	All States	Foliar spray	3 L /ha in min 800 - 1000 L of water	Use weekly spray intervals when conditions favour disease development. To avoid phytotoxicity with some plant species, it is recommended that the product be tested on a few plants of each species prior to the main application.
ORNAMENTALS	Phytophthora Root and Crown rot (<i>Phytophthora</i> spp.)	Qld, NSW, Vic, WA, TAS, NT, ACT	Foliar spray	Knapsack/Boom 170 mL/ 100 L Air Blast 330 mL/100 L	Apply at 4 - 6 weekly intervals when conditions favour disease development. To avoid phytotoxicity with some plant species it is recommended that the product be tested on a few plants of each species prior to the main application. DO NOT apply when ornamental plants are dormant or stressed. DO NOT apply to ornamental plants under extremes of temperature.
PINEAPPLES	Phytophthora Root (<i>Phytophthora cinnamomi</i>) and Heart Rot (<i>Phytophthora parasitica</i>)	Qld, WA	Foliar spray	4 L/ha 1:200 – 1:500 water	Apply to tops, two (2) weeks prior to harvest of planting material.
SUBTERRANEAN CLOVER	Phytophthora Root Rot (<i>Phytophthora clandestina</i>)	NSW, SA VIC, WA, TAS,ACT	Foliar spray	500 mL/ha approx. 200 L water/ha	Apply 8 to 9 days after first irrigation but before second irrigation. Apply in Autumn when Subterranean Clover is at the cotyledon to unifoliate leaf growth stage.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

For effective disease control a good leaf cover must be achieved before a widespread disease outbreak occurs.

CONDITIONS CONDUCTIVE TO DOWNY MILDEW INFECTION

(1) PRIMARY INFECTION - -overnight conditions of:

Temperature	10°C
Rainfall	10 mm
Soil Wetness	24 hrs
Leaf Wetness	3 - 4 hrs at end of a 24 hr period.

(2) SECONDARY INFECTION - Overnight conditions of:

Temperature	11°C
Humidity	98% for at least 4 hours from midnight to dawn
Leaf Wetness	24 hrs

TO AVOID RESISTANT STRAINS OF DOWNY MILDEW DEVELOPING, EZYCROP PHOSPHOROUS SYSTEMIC FUNGICIDE SHOULD BE APPLIED AS CLOSE AS POSSIBLE TO THE DAY OF INFECTION AND ALTERNATED WITH SYSTEMIC FUNGICIDES.

MIXING

Half fill spray vat and with agitation system running, add the required amount of product.

COMPATIBILITY

Compatible with Mancozeb, Copper Oxychloride, Bayleton, Tilt, and most common powdery mildew fungicides. Most foliage nutrients are also compatible. When using in conjunction with 'EC's' always form emulsion with water prior to adding NovaGuard Phosphorous Systemic Fungicide.