EZYCROP PTY LTD
Ezycrop Wetter 1000

APVMA Approval No: 81512/103266

Company Name:

Product Name:





Label Name:	Ezycrop Wetter 1000		
Signal Headings:	READ SAFETY DIRECTIONS BEFORE OPENING OR USING		
Constituent Statements:	ACTIVE CONSTITUENT: 1000 g/L NONIONIC ALCOHOL ETHOXYLATES		
Mode of Action:			
Statement of Claims:	A non-ionic 100% wetting agent, for the improvement of spray coverage when using agricultural chemicals.		
Net Contents:	5 L 20 L 200 L 1000 L		
Restraints:			
Directions for Use:	This section contains file attachment. File Name: Ezy Wetter- DFU - 0715.docx File Size: 23446 bytes		
Other Limitations:			
Withholidng			
Periods:			
Trade Advice:			

General Instructions:

GENERAL INSTRUCTIONS

NOTE: Where product labels recommend the addition of a non-ionic wetting agent as a percentage (%) dilution, add 100 mL Ezycrop Wetter 1000 per 100 litres of spray mixture to give 0.1%, 250 mL to give 0.25% or other rates as required using these examples.

MIXING

For wettable powders and water soluble products, add Ezycrop Wetter 1000 in water before adding the agricultural chemical to the spray tank. For emulsifiable products, add Ezycrop Wetter 1000 after a good emulsion is formed.

Resistance
Warning:

Precautions:

Protections:

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Harmful to fish. DO NOT contaminate ponds, streams, rivers and waterways with chemical or used containers.

Storage and Disposal:

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN.

Store in the closed, original container in a well-ventilated area away from food, drink and stock food. Do not store for prolonged periods in direct sunlight. Store away from oxidising agents. DO NOT re-use container.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Refillable containers: Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage.

Safety Directions:

SAFETY DIRECTIONS

Avoid contact with eyes and skin. When using or preparing product for use, wear overalls, PVC/rubber gloves, goggles and safety shoes. After use and before eating, drinking or smoking wash hands.

First Aid Instructions:

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia: 13 11 26. If swallowed rinse mouth with water. Give several glasses of water to drink. If vomiting occurs, give more water to drink and seek medical advice. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor. If skin contact occurs, remove contaminated clothing and wash thoroughly with soap and water. If irritation occurs seek medical advice.

First Aid Warnings:	

DIRECTIONS FOR USE

Ezycrop Wetter 1000 can be used where the addition of a surfactant is required or recommended on the agricultural chemical label. Variations in conditions of use can require some changes in the amount of Ezycrop Wetter 1000.

SITUATION	RATE/100L	CRITICAL COMMENTS
Non-Selective Herbicides	25 - 100 mL	Use the higher rate on waxy, hairy or hard to
FOLIAR ACTIVE		wet surfaces. Increase rate for dry soils.
Selective Herbicides	25 - 100 mL	Use when a surfactant is recommended on
IN CROP USE		herbicide labels. Use the higher rate on waxy,
		hairy or hard to wet surfaces.
Soil Residual Herbicides	100 - 400 mL	When applied with soil residual herbicides to
		standing weeds, add Ezycrop Wetter 1000 to
		thoroughly wet foliage. Consult product labels
		for individual recommendations.
Horticulture	40 - 100 mL	Vary rate according to volume and nature of
INSECTICIDES, FUNGICIDES		the target.
AND GROWTH REGULATORS		Use only when recommended on product
		labels.
Paraquat and Diquat	25 - 100 mL	Vary rate according to spray volume.
Herbicides		Select rate in accordance with product labels.
Important when grasses are the		
major target		
Pasture Topping	125 mL	
For use with Glyphosate CT		
Watering of Turf and Dust	40 - 100 mL	Use the higher rate when watering turf.
Abatement		
As additive to water		
Fire Fighting	5 mL/100 L	In the field Ezycrop Wetter 1000 has the
Water tanker	2 mL/15 L	advantage of less weight to carry compared
Knapsack		with less concentrated surfactants and of
		being non-flammable. It increases the wetting
		power of water which assists in quick control
		and mopping up of fires. It is of particular value
		where water may not be readily available,
		where a less drenching spray is needed, and a
		tank full will cover a bigger area.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

Ezycrop Wetter 1000
7 July 2015
Directions for Use
Page 1 of 1